



# Expert consensus on neoadjuvant immunotherapy for non-small cell lung cancer

Wenhua Liang<sup>1</sup>, Kaican Cai<sup>2</sup>, Chun Chen<sup>3</sup>, Haiquan Chen<sup>4</sup>, Qixun Chen<sup>5</sup>, Junke Fu<sup>6</sup>, Jian Hu<sup>7</sup>, Tao Jiang<sup>8</sup>, Wenjie Jiao<sup>9</sup>, Shuben Li<sup>1</sup>, Changhong Liu<sup>10</sup>, Deruo Liu<sup>11</sup>, Wei Liu<sup>12</sup>, Yang Liu<sup>13</sup>, Haitao Ma<sup>14</sup>, Xiaojie Pan<sup>15</sup>, Guibin Qiao<sup>16</sup>, Hui Tian<sup>17</sup>, Li Wei<sup>18</sup>, Yi Zhang<sup>19</sup>, Song Zhao<sup>20</sup>, Xiaojing Zhao<sup>21</sup>, Chengzhi Zhou<sup>1</sup>, Yuming Zhu<sup>22</sup>, Ran Zhong<sup>1</sup>, Feng Li<sup>1</sup>, Rafael Rosell<sup>23</sup>, Mariano Provencio<sup>24</sup>, Erminia Massarelli<sup>25</sup>, Mara B. Antonoff<sup>26</sup>, Toyooki Hida<sup>27</sup>, Marc de Perrot<sup>28</sup>, Steven H. Lin<sup>29</sup>, Massimo Di Maio<sup>30</sup>, Antonio Rossi<sup>31</sup>, Dirk De Ruyscher<sup>32</sup>, Robert A. Ramirez<sup>33</sup>, Wolfram C. M. Dempke<sup>34</sup>, D. Ross Camidge<sup>35</sup>, Nicolas Guibert<sup>36</sup>, Raffaele Califano<sup>37</sup>, Qi Wang<sup>38</sup>, Shengxiang Ren<sup>39</sup>, Caicun Zhou<sup>39</sup>, Jianxing He<sup>1</sup>

<sup>1</sup>Department of Thoracic Surgery and Oncology, The First Affiliated Hospital of Guangzhou Medical University, National Center for Respiratory Medicine, State Key Laboratory of Respiratory Disease and National Clinical Research Center for Respiratory Disease, Guangzhou, China; <sup>2</sup>Nanfeng Hospital, Guangzhou, China; <sup>3</sup>Fujian Medical University Union Hospital, Fuzhou, China; <sup>4</sup>Fudan University Shanghai Cancer Center, Shanghai, China; <sup>5</sup>Zhejiang Cancer Hospital, Hangzhou, China; <sup>6</sup>The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China; <sup>7</sup>The First Affiliated Hospital, Zhejiang University, School of Medicine, Hangzhou, China; <sup>8</sup>Tangdu Hospital, Xi'an, China; <sup>9</sup>The Affiliated Hospital of Qingdao University, Qingdao, China; <sup>10</sup>the Second Hospital of Dalian Medical University, Dalian, China; <sup>11</sup>China-Japan Friendship Hospital, Beijing, China; <sup>12</sup>The First Bethune Hospital of Jilin University, Changchun, China; <sup>13</sup>Chinese PLA General Hospital, Beijing, China; <sup>14</sup>The Affiliated Hospital of Soochow University, Suzhou, China; <sup>15</sup>Fujian Provincial Hospital, Fuzhou, China; <sup>16</sup>Guangdong Provincial People's Hospital, Guangzhou, China; <sup>17</sup>Qilu Hospital of Shandong University, Ji'nan, China; <sup>18</sup>Henan Provincial People's Hospital, Zhengzhou, China; <sup>19</sup>Xuanwu Hospital Capital Medical University, Beijing, China; <sup>20</sup>The Affiliated Hospital of Zhengzhou University, Zhengzhou, China; <sup>21</sup>Renji Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China; <sup>22</sup>Shanghai Pulmonary Hospital, Shanghai, China; <sup>23</sup>Department of Medical Oncology, University of Barcelona, Hospital de Badalona Germans Trias i Pujol, Barcelona, Spain; <sup>24</sup>Hospital Universitario Puerta de Hierro-Majadahonda, Madrid, Spain; <sup>25</sup>City of Hope National Cancer Center, Duarte, CA, USA; <sup>26</sup>Department of Thoracic and Cardiovascular Surgery, University of Texas MD Anderson Cancer Center, Houston, TX, USA; <sup>27</sup>Department of Thoracic Oncology, Aichi Cancer Center Hospital, Nagoya, Japan; <sup>28</sup>Toronto Mesothelioma Research Program, Toronto General Hospital, Toronto, ON, Canada; <sup>29</sup>Department of Radiation Oncology, The University of Texas M.D. Anderson Cancer Center, Houston, TX, USA; <sup>30</sup>Department of Oncology, University of Turin/Division of Medical Oncology, Ordine Mauriziano Hospital, Turin, Italy; <sup>31</sup>Oncology Center of Excellence, Therapeutic Science & Strategy Unit, IQVIA, Milan, Italy; <sup>32</sup>Department of Radiation Oncology (MAASTRO), GROW-School for Oncology and Developmental Biology, Maastricht University Medical Centre, Maastricht, The Netherlands; <sup>33</sup>Department of Internal Medicine, Section of Hematology/Oncology, Ochsner Medical Center, Kenner, LA, USA; <sup>34</sup>Department of Hematology and Oncology, University Medical School, Munich, Germany; <sup>35</sup>Division of Medical Oncology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA; <sup>36</sup>Thoracic Oncology Department, Larrey Hospital, University Hospital of Toulouse, Toulouse, France; <sup>37</sup>Department of Medical Oncology, The Christie NHS Foundation Trust, and Division of Cancer Sciences, The University of Manchester, Manchester, UK; <sup>38</sup>Department of Respiratory Medicine, The Second Hospital, Dalian Medical University, Dalian, China; <sup>39</sup>Department of Medical Oncology, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Shanghai, China

*Correspondence to:* Jianxing He. National Center for Respiratory Medicine, China State Key Laboratory of Respiratory Disease and National Clinical Research Center for Respiratory Disease, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China. Email: drjianxing.he@gmail.com.

Submitted Oct 21, 2020. Accepted for publication Dec 24, 2020.

doi: 10.21037/tlcr-2020-63

View this article at: <http://dx.doi.org/10.21037/tlcr-2020-63>

## Introduction

Lung cancer is the leading cause of cancer-related death worldwide and in China (1). According to the statistics of the National Cancer Center of China, there were 733,300 new cases of non-small cell lung cancer (NSCLC) and approximately 610,200 related deaths in 2015 (2). For patients with early staged disease, surgery is the mainstay of treatment, and it is commonly followed by adjuvant chemotherapy for patients with locally advanced resectable NSCLC. Although complete surgical resection may be curative for NSCLC, 25–70% of patients (with different proportion according to stage) eventually relapse despite complete resection (3). Platinum-based adjuvant chemotherapy has been shown to marginally increase the 5-year survival rate of patients by 4–8% (4–6). Even after treatment with surgery and indicated adjuvant therapies in eligible cases, approximately 20–30% of stage I, 50% of stage II, and 60% of stage IIIA patients still die within 5 years (7). In the past decade, experts have conducted a number of investigations on the perioperative management of resectable NSCLC; however, progress remains slow, and patients still have a high risk of recurrence and death.

Neoadjuvant therapy is defined as any therapy delivered prior to definitive local therapy intended to increase the cure rate. It provides several theoretical benefits in managing such patients with NSCLC. In the setting of, neoadjuvant therapy given prior to radical surgery this approach can also have the goals of downstaging, improving the resection rate, and more promptly treating subclinical micro-metastases than adjuvant approaches, delivered after the definitive local therapy. In addition, the compliance with neoadjuvant therapy has been shown to be better than in the adjuvant setting, and the biological effect of the neoadjuvant therapy can be analyzed directly in the resected tumor specimens (8). A meta-analysis on patients with stage IB–IIIA NSCLC that compared chemotherapy plus subsequent surgery *vs.* surgery alone showed that the 5-year survival rate was 5% higher after receiving neoadjuvant chemotherapy (NCT) (9). Therefore, the comprehensive NSCLC data suggest that, for resectable NSCLC, NCT improves survival compared with surgery alone but appear to show no significant survival benefit compared with adjuvant chemotherapy (10).

In the last 5 years, immune checkpoint inhibitors (ICIs) have profoundly changed the treatment paradigm for patients with advanced NSCLC (11–15). Immunotherapy has provided hope for long-term survival benefits to

a minority of patients with metastatic lung cancer. For treatment-naïve patients with driver mutation-negative NSCLC, the 5-year survival rate of single agent pembrolizumab was 23.2%; for the previously treated patients with driver mutation-negative NSCLC, the 5-year survival rates of single agent pembrolizumab and nivolumab were 15.5% and 16%, respectively (16,17). Given the profound impact made by immunotherapy drugs for patients with advanced disease, significant attention has been directed in recent years toward investigating the potential role for early-stage NSCLC patients, and whether they, too, can achieve long-term benefits from the inclusion of immunotherapy into their treatment algorithms.

Many phase Ib/II clinical trials have reported promising results, and a series of large-scale phase III clinical trials are underway. However, these various investigations have employed different strategies of neoadjuvant immunotherapy, in terms of the specific regimens as well as number of treatment cycles (18). To better guide Chinese thoracic surgeons in the neoadjuvant immunotherapy of NSCLC, well-known thoracic surgeons in China participated in an in-depth discussion on the hot topics and controversial issues of neoadjuvant immunotherapy and formed the *Expert consensus on neoadjuvant immunotherapy for non-small-cell lung cancer* by incorporating the latest evidence on neoadjuvant immunotherapy.

### **Consensus 1: preoperative use of neoadjuvant immunotherapy with or without platinum-based chemotherapy for patients with resectable stage IB–IIIA NSCLC may be considered**

In 1994, Rosell and Roth published two classic prospective randomized controlled trials of NCT for stage IIIA NSCLC, and since then a number of studies since have demonstrated that, for early-stage NSCLC, the efficacy of preoperative NCT plus surgery is similar to that of surgery plus postoperative adjuvant chemotherapy, and both strategies including systemic treatment are better than surgery alone in terms of overall survival (OS) (9,19–21). Due to the relevant toxicity of neoadjuvant/adjuvant chemotherapy, patient tolerance can, at times, be poor. Compared with surgery alone, the addition of neoadjuvant or adjuvant chemotherapy only modestly improves the 5-year OS rate by about 5% (9,21). Thus, there is a clear need for well-tolerated and effective neoadjuvant and adjuvant therapies for resectable NSCLC. Compared with conventional chemotherapy, immunotherapy is better

tolerated in most patients, and treatment-related toxicities may have minimal influence on ability to complete surgical resection. Immunotherapy is being actively explored for the perioperative management of NSCLC.

When programmed death 1 (PD-1)/programmed death-ligand 1 (PD-L1) inhibitors kill tumors, antigen-presenting cells (APC) are needed so that the tumor cells can be recognized by host T cells. By blocking the PD-1/PD-L1 interaction with inhibitory antibodies, immunotherapy allows the activated T cells to release cytokines, perforins, granzymes and others to kill tumor cells. In terms of mechanisms involved, when the tumor volume is relatively large, APC bears a relatively large antigen load, which elicits a stronger antitumor T cell response. Therefore, neoadjuvant immunotherapy could theoretically be better than adjuvant immunotherapy and may yield an even greater survival long-term benefit (22). Preclinical studies support these hypotheses. In animal models, compared with adjuvant immunotherapy, neoadjuvant immunotherapy resulted in greater prolongation of median survival time and a higher survival rate (23-25).

No data from large-scale phase III clinical trials of neoadjuvant immunotherapy are yet available, but the results of several phase II clinical trials showed that neoadjuvant immunotherapy may play an important role in the multi-disciplinary management of early-stage NSCLC (26-31). The results of trials of neoadjuvant immunotherapy were summarized in *Table 1*.

CheckMate-159 is a phase II clinical trial which evaluated the safety and feasibility of a preoperative neoadjuvant immunotherapy with two-cycle nivolumab monotherapy, in resectable stage I-III NSCLC. This single-arm study enrolled 22 patients, of which 20 patients received two cycles of the experimental treatment and 20 underwent complete surgical resection. This study showed that neoadjuvant nivolumab monotherapy was safe, tolerable and effective. The major pathologic response (MPR, defined as residual viable tumor cells less than 10%) rate was 45% (26). The phase II clinical trial, TOP1501, was designed to evaluate the safety and feasibility of another preoperative neoadjuvant immunotherapy, two cycles of pembrolizumab monotherapy, in patients with stage IB-III NSCLC. This monotherapy was also found to be safe and efficacious. The MPR rate was 28%, and up to 80% of patients had pathological remission  $\geq 50\%$  (28). The LCMC3 phase II clinical study of PD-L1 inhibitors showed that in patients with stage IB-III NSCLC (plus selected stage IIIB patients), two-cycle

atezolizumab monotherapy was found to be well-tolerated and have promising outcomes. The MPR rate was 19%, the pathologic complete remission (pCR, defined as no residual viable tumor) was 5%, and the MPR was unrelated to PD-L1 expression (27). In addition to the above immunotherapies, neoadjuvant immunotherapy combined with chemotherapy is associated with higher MPRs in cross-trial comparisons and a suggestion of longer OS utilizing the same cross-trial comparators. A phase II study by Shu *et al.* (32) aimed to investigate the safety and efficacy of preoperative neoadjuvant immunotherapy in the form of four cycles of atezolizumab monotherapy plus carboplatin plus albumin-bound paclitaxel in patients with stage IB-III NSCLC, and the results revealed that this treatment algorithm was feasible, tolerable, and beneficial. The MPR was 50%, and the pCR reached 21.4%. The NADIM study presented at ASCO in 2019 aimed to assess the safety and efficacy of preoperative neoadjuvant immunotherapy in the form of three-cycle nivolumab monotherapy plus carboplatin plus paclitaxel in patients with stage IIIA (N2 or T4N0) NSCLC, and results were promising. The MPR was 83%, the pCR was 71%, and tumor was down-staged in 90% of the patients (33). Currently, the experimental groups of all phase III clinical trials are including neoadjuvant immunotherapy with PD-1/PD-L1 inhibitors plus platinum-based doublet chemotherapy, such as CheckMate-816 (NCT02998528), KEYNOTE-671 (NCT03425643), IMpower-030 (NCT03456063), AEGEAN (NCT03800134), CheckMate-77T (NCT04025879). The ongoing trials were summarized in *Table S1*.

In summary, this consensus suggests that preoperative use of neoadjuvant immunotherapy with or without platinum-based chemotherapy might be considered for patients with resectable stage IB-III NSCLC.

**Consensus 2: there is no evidence that molecular markers uniformly predict the efficacy of neoadjuvant immunotherapy so that biomarker-based selection is not essential. However, it should be cautious to use neoadjuvant single agent immunotherapy in patients with potentially negative factors, such as epidermal growth factor receptor (EGFR)—sensitive mutation/ALK fusion**

PD-1/PD-L1 inhibitors have changed the treatment strategies of advanced NSCLC, but we know that not all

**Table 1** The results of trials of neoadjuvant immunotherapy

Study title	NCT.no	Key inclusion and exclusion criteria	Drug	Major result
A Randomized, Double-Blind, Phase III Study of Platinum + Pemetrexed Chemotherapy With or Without Pembrolizumab (MK-3475) in First Line Metastatic Non-squamous Non-small Cell Lung Cancer Subjects (KEYNOTE-189)	NCT02578680	Patients with previously untreated metastatic nonsquamous NSCLC without EGFR or ALK mutations	Cisplatin; carboplatin; pemetrexed; dexamethasone; pembrolizumab	(I) OS at 12 months was 69.2% in the pembrolizumab-combination group versus 49.4% in the placebo-combination group. (II) PFS was 8.8 months in the pembrolizumab-combination group and 4.9 months in the placebo-combination group. (III) Adverse events of grade 3 or higher occurred in 67.2% of the patients in the pembrolizumab-combination group and in 65.8% of those in the placebo-combination group
A Randomized, Double-Blind, Phase III Study of Carboplatin-Paclitaxel/ Nab-Paclitaxel Chemotherapy With or Without Pembrolizumab (MK-3475) in First Line Metastatic Squamous Non-small Cell Lung Cancer Subjects (KEYNOTE-407)	NCT02775435	Untreated metastatic squamous NSCLC	Paclitaxel; nab-paclitaxel; carboplatin; pembrolizumab	(I) OS was 15.9 months in the pembrolizumab-combination group and 11.3 months in the placebo-combination group. (II) PFS was 6.4 months in the pembrolizumab-combination group and 4.8 months in the placebo-combination group. (III) Adverse events of grade 3 or higher occurred in 69.8% of the patients in the pembrolizumab-combination group and in 68.2% of the patients in the placebo-combination group
A Randomized, Open Label, Phase III Study of Overall Survival Comparing Pembrolizumab (MK-3475) Versus Platinum Based Chemotherapy in Treatment Naïve Subjects With PD-L1 Positive Advanced or Metastatic Non-Small Cell Lung Cancer (Keynote 042)	NCT02220894	Patients with previously untreated metastatic NSCLC without EGFR or ALK mutations and with an ECOG performance status score of 0 or 1, life expectancy 3 months or longer, and a PD-L1 TPS of 1% or greater.	Carboplatin; paclitaxel; pemetrexed; pembrolizumab	(I) The median survival values by TPS population ( $\geq 50\%$ , $\geq 20\%$ , $\geq 1\%$ ) were 20.0 months for pembrolizumab versus 12.2 months for chemotherapy, 17.7 months versus 13.0 months, and 16.7 months versus 12.1 months, respectively. (II) Treatment-related adverse events of grade 3 or worse occurred in 18% of the patients in the pembrolizumab group and in 41% of those in the chemotherapy group and led to death in 2% and 2% patients, respectively
Phase I Study of Single Agent Pembrolizumab (MK-3475) in Patients With Progressive Locally Advanced or Metastatic Carcinoma, Melanoma, and Non-Small Cell Lung Carcinoma (KEYNOTE 001)	NCT01295827	Patients had confirmed locally advanced/metastatic NSCLC and provided a contemporaneous tumor sample for PD-L1 evaluation by immunohistochemistry using the 22C3 antibody.	Pembrolizumab	(I) Median OS was 22.3 months in treatment-naïve patients and 10.5 months in previously treated patients. (II) Estimated 5-year OS was 23.2% for treatment-naïve patients and 15.5% for previously treated patients. (III) In patients with a PD-L1 tumor proportion score of 50% or greater, 5-year OS was 29.6% and 25.0% in treatment-naïve and previously treated patients, respectively. (IV) Compared with analysis at 3 years, only three new-onset treatment-related grade 3 adverse events occurred
A Phase 1, Open-Label, Multicenter, Multidose, Dose Escalation Study of BMS-936558 (Nivolumab) in Subjects With Selected Advanced or Recurrent Malignancies	NCT00730639	Advanced melanoma, RCC, or NSCLC who received nivolumab and were enrolled between October 30, 2008, and December 28, 2011	Nivolumab	OS was significantly longer among patients with treatment-related AEs of any grade (median, 19.8 months) or grade 3 or more (median, 20.3 months) compared with those without treatment-related AEs (median, 5.8 months)

**Table 1** (continued)

Table 1 (continued)

Study title	NCT.no	Key inclusion and exclusion criteria	Drug	Major result
Neoadjuvant Nivolumab, or Nivolumab in Combination With Ipilimumab, in Resectable Non-Small-Cell Lung Cancer.	NCT02259621	Adults with untreated, surgically resectable early (stage I, II, or IIIA) NSCLC	Nivolumab	Neoadjuvant nivolumab was associated with few side effects, did not delay surgery, and induced a major pathological response in 45% of resected tumors
A Randomized Open-Label Phase III Trial of MK-3475 Versus Platinum Based Chemotherapy in 1L Subjects With PD-L1 Strong Metastatic Non-Small Cell Lung Cancer	NCT02142738	Previously untreated advanced NSCLC with PD-L1 expression on at least 50% of tumor cells and without EGFR or ALK mutations	Paclitaxel; carboplatin; pemetrexed; cisplatin; gemcitabine; pembrolizumab	(I) PFS was 10.3 months in the pembrolizumab group versus 6.0 months in the chemotherapy group. (II) OS at 6 months was 80.2% in the pembrolizumab group versus 72.4% in the chemotherapy group. (III) The response rate was higher in the pembrolizumab group than in the chemotherapy group (44.8% vs. 27.8%). (IV) Treatment-related adverse events of any grade were less frequent (occurring in 73.4% vs. 90.0% of patients), as were grade 3, 4, or 5 treatment-related adverse events (26.6% vs. 53.3%)
An Open-Label, Randomized Phase 3 Trial of Nivolumab, or Nivolumab Plus Ipilimumab, or Nivolumab Plus Platinum Doublet Chemotherapy Versus Platinum Doublet Chemotherapy-Naive Stage IV or Recurrent Non-Small Cell Lung Cancer (NSCLC)	NCT02477826	Patients with stage IV or recurrent NSCLC that was not previously treated	Nivolumab; ipilimumab; carboplatin; cisplatin; gemcitabine; pemetrexed; paclitaxel	(I) The 1-year PFS rate was 42.6% with nivolumab plus ipilimumab versus 13.2% with chemotherapy, and the median PFS was 7.2 months versus 5.5 months. (II) The objective response rate was 45.3% with nivolumab plus ipilimumab and 26.9% with chemotherapy. (III) The rate of grade 3 or 4 treatment-related adverse events was 31.2% with nivolumab plus ipilimumab and 36.1% with chemotherapy
An Open-Label Randomized Phase III Trial of BMS-936558 (Nivolumab) Versus Docetaxel in Previously Treated Metastatic Non-squamous Non-small Cell Lung Cancer (NSCLC)	NCT01673867	Patients with nonsquamous NSCLC that had progressed during or after platinum-based doublet chemotherapy	Nivolumab; docetaxel	(I) The OS was 12.2 months in the nivolumab group and 9.4 months in the docetaxel group. At 1 year, the OS rate was 51% with nivolumab versus 39% with docetaxel. OS rate at 18 months was 39% with nivolumab versus 23% with docetaxel. (II) The response rate was 19% with nivolumab versus 12% with docetaxel. (III) The rate of PFS at 1 year was higher with nivolumab than with docetaxel (19% and 8%, respectively). (IV) Treatment-related adverse events of grade 3 or 4 were reported in 10% of the patients in the nivolumab group, as compared with 54% of those in the docetaxel group.

Table 1 (continued)

Table 1 (continued)

Study title	NCT.no	Key inclusion and exclusion criteria	Drug	Major result
A Phase II/III Randomized Trial of Two Doses of MK-3475 (SCH900475) Versus Docetaxel in Previously Treated Subjects With Non-Small Cell Lung Cancer	NCT01905657	Patients with previously treated non-small-cell lung cancer with PD-L1 expression on at least 1% of tumour cells	Pembrolizumab; docetaxel	(I) OS was 10.4 months with pembrolizumab 2 mg/kg, 12.7 months with pembrolizumab 10 mg/kg, and 8.5 months with docetaxel. (II) PFS was 3.9 months with pembrolizumab 2 mg/kg, 4.0 months with pembrolizumab 10 mg/kg, and 4.0 months with docetaxel (III) Among patients with at least 50% of tumour cells expressing PD-L1, OS was significantly longer with pembrolizumab 2 mg/kg than with docetaxel (14.9 vs. 8.2 months) and with pembrolizumab 10 mg/kg than with docetaxel (17.3 vs. 8.2 months). PFS was significantly longer with pembrolizumab 2 mg/kg than with docetaxel (5.0 vs. 4.1 months) and with pembrolizumab 10 mg/kg than with docetaxel (5.2 vs. 4.1 months) (IV) Grade 3-5 treatment-related adverse events were less common with pembrolizumab than with docetaxel (13% of the patients given 2 mg/kg, 16% of given 10 mg/kg, and 35% of 309 given docetaxel)
A Randomized, Open Label, Phase III Study of Overall Survival Comparing Pembrolizumab (MK-3475) Versus Platinum Based Chemotherapy in Treatment Naïve Subjects With PD-L1 Positive Advanced or Metastatic Non-Small Cell Lung Cancer (Keynote 042)-China Extension Study	NCT03850444	Chinese participants with PD-L1-positive NSCLC	Pembrolizumab; carboplatin; paclitaxel; pemetrexed	(I) OS (TPS of $\geq 50\%$ ) was 20.0 months in the pembrolizumab group and 14.0 months in the Chemotherapy group. OS (TPS of $\geq 20\%$ ) was 20.0 months in the pembrolizumab group and 13.7 months in the Chemotherapy group. OS (TPS of $\geq 1\%$ ) was 20.0 months in the pembrolizumab group and 13.7 months in the Chemotherapy group. (II) PFS (TPS of $\geq 50\%$ ) was 8.3 months in the pembrolizumab group and 6.5 months in the Chemotherapy group. PFS (TPS of $\geq 20\%$ ) was 6.3 months in the pembrolizumab group and 6.5 months in the Chemotherapy group. PFS (TPS of $\geq 1\%$ ) was 6.3 months in the pembrolizumab group and 6.4 months in the Chemotherapy group
A Randomized, Double-Blind, Phase III Study of Platinum + Pemetrexed Chemotherapy With or Without Pembrolizumab (MK-3475) in First Line Metastatic Non-squamous Non-small Cell Lung Cancer Subjects (KEYNOTE-189)	NCT03950674	Adult Japanese participants with advanced or metastatic nonsquamous NSCLC who have not previously received systemic therapy for advanced disease	Dexamethasone; cisplatin; carboplatin; pemetrexed	(I) PFS was 16.5 months in the pembrolizumab group and 7.1 months in the Chemotherapy group. (II) ORR was 56% in the pembrolizumab group and 33.3% in the Chemotherapy group

Table 1 (continued)

Table 1 (continued)

Study title	NCT.no	Key inclusion and exclusion criteria	Drug	Major result
An Open-label Randomized Multinational Phase 3 Trial of Nivolumab Versus Docetaxel in Previously Treated Subjects With Advanced or Metastatic Non-small Cell Lung Cancer (CheckMate 078: CHECK point Pathway and nivolumab Clinical Trial Evaluation 078)	NCT02613507	Advanced or Metastatic NSCLC who have failed prior platinum-based doublet chemotherapy	Nivolumab; docetaxel	(I) OS was 11.99 months in the Nivolumab group and 9.63 months in the Docetaxel group. (II) ORR was 16.6% in the Nivolumab group and 4.2% in the Docetaxel group
A Phase 3, Randomized Study of Nivolumab Plus Ipilimumab in Combination With Chemotherapy vs. Chemotherapy Alone as First Line Therapy in Stage IV Non-Small Cell Lung Cancer	NCT03215706	Stage IV NSCLC have not previously received systemic therapy	Ipilimumab; Nivolumab; Carboplatin; Paclitaxel; Pemetrexed; Cisplatin	(I) OS was 14.13 months in the Nivolumab + Ipilimumab + Chemotherapy group and 10.74 months in the Chemotherapy group. (II) PFS was 6.83 months in the Nivolumab + Ipilimumab + Chemotherapy group and 4.96 months in the Chemotherapy group. (III) ORR was 37.7% in the Nivolumab + Ipilimumab + Chemotherapy group and 25.1% in the Chemotherapy group
An Open-Label, Randomized, Phase 3 Trial of Nivolumab Versus Investigator's Choice Chemotherapy as First-Line Therapy for Stage IV or Recurrent PD-L1+ Non-Small Cell Lung Cancer	NCT02041533	Stage IV or Recurrent PD-L1+ NSCLC	Nivolumab; Gemcitabine; Cisplatin; Carboplatin; Paclitaxel; Pemetrexed	(I) PFS was 4.21 months in the Nivolumab group and 5.82 months in the Chemotherapy group. PFS (TPS of $\geq 5\%$ ) was 4.21 months in the Nivolumab group and 5.88 months in the Chemotherapy group. (II) OS was 13.73 months in the pembrolizumab group and 13.80 months in the Chemotherapy group. OS (TPS of $\geq 5\%$ ) was 14.36 months in the pembrolizumab group and 13.21 months in the Chemotherapy group
A Dose Frequency Optimization, Phase IIIb/IV Trial of Nivolumab 240 mg Every 2 Weeks vs. Nivolumab 480 mg Every 4 Weeks in Subjects With Advanced or Metastatic Non-small Cell Lung Cancer Who Received up to 12 Months of Nivolumab at 3 mg/kg or 240 mg Every 2 Weeks	NCT02713867	Advanced/metastatic (Stage IIIb/IV) NSCLC	Nivolumab	(I) PFSR at 6 months was 0.76 in the Nivolumab 480mg Q4W group and 0.79 in the Nivolumab 240 mg Q2W group. (II) OS was at 6 months 0.966 of 180 patients in the Nivolumab 480 mg Q4W group and 0.956 183 patients in the Nivolumab 240 mg Q2W group

patients can benefit from them. For patients with driver mutation–negative NSCLC, the response rate of PD-1/PD-L1 inhibitor monotherapy has been approximately 14–20%, and that of first-line monotherapy in advanced NSCLC patients with PD-L1 expression  $\geq 50\%$  is approximately 50% (34). Therefore, it is critical to identify predictive biomarkers to select patients who can benefit from PD-1/PD-L1 inhibitors, avoiding the treatment to those patients who will not benefit. The most promising marker as such is PD-L1 so far. Supported by well-founded evidence. KEYNOTE-024 and KEYNOTE-042 studies indicated that among patients with high or positive PD-L1 expression, pembrolizumab monotherapy was superior to the standard chemotherapy. The higher the PD-L1 expression, the more benefit the patients received from single agent immunotherapy. Therefore, all guidelines consistently recommend PD-L1 as the companion diagnostic for patients with driver mutation–negative NSCLC receiving immunotherapy (13,35). The IMpower110 study also showed that among the patients with driver mutation–negative NSCLC and high PD-L1 expression in tumor cells and/or interstitial cells (TC3/IC3), atezolizumab monotherapy was superior to the standard chemotherapy (36). Some results of KEYNOTE-016, KEYNOTE-164, KEYNOTE-1022, KEYNOTE-028, KEYNOTE-177 and KEYNOTE-158 indicated that the objective remission rate of pembrolizumab monotherapy reached 39.6% in solid-tumor patients with microsatellite instability high (MSI-H) or mismatch repair deficiency. Based on this finding, the United States Food and Drug Administration (FDA) approved pembrolizumab monotherapy for patients with disease progression after previous treatment or metastatic solid tumor patients with MSI-H or mismatch repair deficiency, which is the first indication for cancer treatment based on biomarkers (37). However, these features are very rare in NSCLC.

The use of PD-L1 as a biomarker, however, remains to be complicated by a number of factors including the variability in tissue collection timing, the antibody and methodology used for staining (including the definition of positivity and the non-standardised test design), the heterogeneity and dynamic of PD-L1 expression within different tumors, and the role of PD-L1 expression on tumor-infiltrating lymphocytes and other immune cells versus the malignant cell population. In addition, PD-L1 is regarded to be a biological continuum and therefore might be of limited value as a biomarker in this subset of patients. Pembrolizumab (anti-PD-1) is approved (EMA, FDA) for

first-line treatment of NSCLC patients with advanced or metastatic cancers (with PD-L1 expression  $\geq 50\%$  using the Dako 22C3 IHC assay), whereas for second-line treatment a PD-L1 expression of  $\geq 1\%$  is required. Both, nivolumab (anti-PD-1) and atezolizumab (anti-PD-L1) are also approved (EMA, FDA) in the first- and second-line setting, but PD-L1 screening is only mandatory for atezolizumab, however, complementary PD-L1 diagnostics are approved for NSCLC. Durvalumab (anti-PD-L1) is currently for stage non-resectable IIIA/IIIB NSCLC, PD-L1 testing is recommended. To date, many groups of oncologists are attempting to establish better predictive biomarkers in NSCLC for monoclonal antibodies targeting the PD-1/PD-L1 axis to select patients who might have a greater benefit from immune checkpoint therapies.

In addition to PD-L1 and MSI-H, tumor mutation burden (TMB) is also considered a potential biomarker for immunotherapy. Noteworthy, TMB can be assessed in different ways and, as a continuous variable, ‘high’ can also be variably defined. Although the retrospective analyses of KEYNOTE-042 and KEYNOTE-189 trials, and the prospective study CheckMate-227 all showed that TMB could not effectively predict the efficacy of immunotherapy alone or immunotherapy combined with chemotherapy as first-line treatment of advanced NSCLC (38), the KEYNOTE-158 results showed that, among patients with disease progression or metastases after previous treatment, the outcomes of single-agent pembrolizumab in those with high TMB were significantly better than in those with low TMB. Based on this, the FDA approved the pembrolizumab monotherapy for patients with high TMB as defined by  $\geq 10$  mutations/Mb and disease progression after previous treatment and for metastatic solid tumor patients (39).

The IMpower150 trial is the first phase III study to demonstrate a clinically meaningful and significant PFS benefit with atezolizumab plus bevacizumab and chemotherapy (paclitaxel and carboplatin) versus bevacizumab plus chemotherapy in the first-line setting of advanced or metastatic NSCLC (8.3 versus 6.8 months, HR =0.62,  $P < 0.0001$ ) and the PFS benefit was seen regardless of the PD-L1 status in all patients. The PFS benefit, however, was even more pronounced in patients expressing a T effector gene signature (11.3 versus 6.8 months, HR =0.51,  $P < 0.0001$ ) indicating that the expression of gene signatures may be more robust to predict clinical response following treatment with PD-L1 inhibitors (40).

Of note, in experimental systems with NSCLC-bearing xenografts, the tumor microenvironment (TME) has also

been found to significantly contribute to the response to CPIs. Several lines of evidence have suggested that tumor-associated macrophages (TAMs: positivity for PD-L1, CD133, and CD163) within the TME can reduce responses to immunotherapies by suppressing CD8-positive cytotoxic T cells (CTLs). They are also regarded to be responsible for the hyperprogression following immunotherapy (41). In addition, cancer-associated fibroblasts (CAFs) have also been shown to suppress responses to CPIs by decreasing CD8-positive CTLs. Ford *et al.* (42) have provided the first evidence that TGF- $\beta$  can stimulate NADPH-Oxidase 4 (NOX-4) which in turn produces reactive oxygen species that can activate CAFs and thereby confer resistance to IO treatments. Although so far these are experimental approaches, it is likely that the CAF and TAM status will guide immunotherapy treatment strategies in the neoadjuvant setting in the future as well.

Differently from advanced NSCLC, immunotherapy for early-stage NSCLC is still under investigation. Clinical studies have yet to find biomarkers that can predict the efficacy of neoadjuvant immunotherapy (27,43). The NEOSTAR study showed that, when using neoadjuvant nivolumab monotherapy or nivolumab combined with ipilimumab monotherapy to treat stage I–IIIA NSCLC, the baseline PD-L1 expression in patients obtaining complete/partial response (CR/PR) was higher than that in patients obtaining stable/progressive disease (SD/PD), and the baseline PD-L1 expression in patients obtaining MPR was higher than those who did not (43). In the LCMC3 study, neoadjuvant atezolizumab monotherapy was given to patients with stage IB–IIIB NSCLC. Pathological remission and MPR were observed regardless of the expression level of PD-L1, and TMB had no correlation with pathological remission and MPR (27). Immunotherapy for early-stage NSCLC is still under investigation, and biomarkers in this setting are also being explored. All currently completed studies are small-scale phase I/II clinical trials, and the role of biomarkers, including PD-L1, MSI-H, T<sub>eff</sub> cell signatures, and TMB, remains unclear, warranting further exploration.

Over the past 20 years, with the discovery of driver genes such as *EGFR*, *ROS*, *RET*, and *ALK* and the development of targeted agents, the treatment of NSCLC has entered the era of precision drug therapy, greatly improving the prognosis of molecularly selected subgroups of patients with advanced NSCLC (44). As described above, ICIs can bring long-term benefits to patients with driver mutation-negative NSCLC (11–13). Therefore, many researchers

have explored whether ICIs can also bring long-term benefits to NSCLC patients with driver mutations.

CheckMate-057 is a phase III clinical study aiming to explore the efficacy of nivolumab monotherapy as the second-line treatment of advanced nonsquamous NSCLC, and its efficacy was compared to that of docetaxel. In this study, 82 *EGFR*-mutated NSCLC patients and 21 patients with *ALK* translocation were included, and they all had previously received the first-line platinum-based doublet chemotherapy. Subgroup analysis of *EGFR*-mutated NSCLC patients showed that OS and progression-free survival (PFS) were not improved in patients receiving nivolumab monotherapy compared to those receiving docetaxel (45). Another III randomized clinical trial, KEYNOTE-010, comparing pembrolizumab *vs.* docetaxel, performed subgroup analysis on *EGFR*-mutated NSCLC patients and showed that pembrolizumab did not improve OS compared to docetaxel (46). A recent, real-world, retrospective analysis showed that *EGFR*-mutated and *ALK*-rearranged NSCLC patients did not benefit from PD-1/PD-L1 monotherapy. A number of clinical trials and retrospective studies have shown that for advanced *EGFR*-mutated or *ALK*-rearranged patients, the chance of benefit with immunotherapy is modest, and the addition of ICIs on the corresponding targeted therapies does not bring an additional benefit but increases the occurrence of toxic and side effects. Further support for this proposal came from the observation that *EGFR* mutations may decrease PD-L1 expression (47).

In addition to the poor efficacy of immunotherapy in NSCLC patients with driver mutations (positive *EGFR* and *ALK*), genes such as *STK11* that are tumor suppressor genes are frequently mutated in NSCLC and may reduce cluster of differentiation (CD) 8<sup>+</sup> T cell density or function, thus affecting the tumor-related immune response through multiple pathways (48). The MYSTIC study showed that the prognosis of patients with *STK11*-mutated NSCLC receiving single-agent durvalumab was poor (49). The exploratory analysis of KEYNOTE-042 showed that regardless of the *STK11* mutation status, the patients who received pembrolizumab monotherapy showed better PFS and OS than the patients who received standard chemotherapy (50). However, complete responses to nivolumab in patients with *STK11* mutation have been also reported (51). At present, it is not clear whether *STK11* mutation is a prognostic factor or a predictive factor in patients with NSCLC receiving for PD-1/PD-L1 inhibitor therapy.

Although the roles of driver genes, such as *EGFR* and *ALK*, and inhibitory genes, such as *STK11*, in immunotherapy for early-stage NSCLC are still not clear. Based on the above data, patients with potentially unfavorable factors such as EGFR-sensitive mutation or ALK fusion should be cautious to receive neoadjuvant single agent immunotherapy.

**Consensus 3: for neoadjuvant immunotherapy, two to four cycles are recommended, and after every two cycles, review and evaluation should be performed to update the treatment plan**

The purpose of neoadjuvant immunotherapy is to enhance downstaging, improve the R0 resection rate, and subsequently treat subclinical micrometastases. A short course of neoadjuvant immunotherapy may not be adequate for immunotherapy to have an effect, but if the duration of immunotherapy is too long, tumor progression may lead to the loss of surgical window of opportunity, so the length of neoadjuvant immunotherapy is very important (52). A preclinical study showed that the presence of a primary tumor appeared to be key to the efficacy of neoadjuvant immunotherapy, and the efficacy was closely correlated with the timing of tumor resection after neoadjuvant immunotherapy (53). To prevent progression in patients with drug resistance, the International Neoadjuvant Melanoma Consortium (INMC) recommends six to eight weeks of neoadjuvant therapy for melanoma, depending on the cycle length of the clinical trial (54). At present, the effects of neoadjuvant immunotherapy on early-stage NSCLC are known only from phase I/II clinical trials, without large-scale phase III trials. The neoadjuvant single agent immunotherapy in CheckMate159, LCMC3, and TOP1501, was performed for two cycles, and surgery was performed 28–56 days after the first cycle (26–28). Neoadjuvant immunotherapy combined with chemotherapy (NADIM: phase II with N=51) or combination of two checkpoint inhibitors (PD-1 and CTLA-4) (NEOSTAR: phase II with N=44) was performed for three to four cycles, and surgery was performed 3–7 weeks after the end of the neoadjuvant therapy (43). Currently, the role of neoadjuvant immunotherapy in early-stage NSCLC is still not fully elucidated, and review after every two cycles is recommended to assess tumor remission and update the treatment plan. However, as there is a low predictive value of CT scans or of RECIST criteria, in case no distant metastases are found, patients should proceed to surgical

resection if still feasible.

In summary, two to four cycles are recommended for neoadjuvant immunotherapy, and re-evaluate after every two cycles is recommended to refine the treatment plan.

**Consensus 4: the benefit from neoadjuvant immunotherapy should be preferably assessed by positron-emission tomography (PET)-computed tomography (CT), in conjunction with serum tumor markers and/or circulating tumor DNA (ctDNA) load**

Tumor shrinkage after the treatment is clear evidence for the antitumor activity of neoadjuvant therapy, and the objective remission rate is an important indicator to evaluate tumor shrinkage and antitumor activity. CT is often used to assess the response of NSCLC patients to neoadjuvant therapy. The Response Evaluation Criteria in Solid Tumors (RECIST) by CT is an important predictor of OS in NSCLC patients after NCT (55). However, the histopathological response of 41–45% patients may be inconsistent with the CT evaluation (26,56). Changes in inflammation and interstitial or fibrotic components of tumors may affect the CT results, leading to the inability of CT imaging to accurately predict histopathological responses after neoadjuvant therapy. When using imaging examinations, the efficacy is determined by continuously measuring changes in tumor size in these patients, which has inherent limitations. Therefore, conventional imaging combined with metabolic imaging may be required to determine efficacy (56). Some investigators have suggested that PET-CT may be more advantageous in the evaluation of neoadjuvant therapy because the uptake of <sup>18</sup>F-fluorodeoxyglucose by the tumor is closely related to the proliferative activity and the number of remaining vital tumor cells. Multiple PET-CT studies have shown that PET-CT can effectively evaluate the efficacy of neoadjuvant therapy. Although the response to neoadjuvant therapy as seen on PET-CT may be related to an improved prognosis, the role of PET-CT in neoadjuvant therapy at early stage still requires further exploration (31,55,57). In addition, incorporating serum tumor markers or ctDNA may increase the accuracy of tumor load assessment (58,59). This is particularly important for assessment of neoadjuvant immunotherapy, given that discrepancy between radiologic and pathologic response, as well as pseudo progression, have been extensively reported (24,28).

In summary, the benefit of neoadjuvant immunotherapy is preferred to be assessed by PET-CT, in conjunction with tumor markers and/or ctDNA load. However, more research on the accuracy and the need for standardization, especially for ctDNA measurements, of these modalities is needed before they can be introduced in standard of care.

#### **Consensus 5: surgery can be performed 4–6 weeks after the last cycle of neoadjuvant immunotherapy**

It is very difficult to determine the timing of surgery after neoadjuvant immunotherapy. For neoadjuvant chemotherapy, early surgery may lead to serious surgical complications, while delayed surgery may lead to tumor progression (52). However, neoadjuvant immunotherapy is distinct from neoadjuvant chemotherapy in terms of adverse events. Before determining the optimal time for neoadjuvant immunotherapy and surgery, it is important to understand the T cell amplification cycle, determine the best time for effector cells to exert their effects, and time the tumor resection for when the impact on antitumor immunity will be the least. Experimentally, this may be very challenging, though recent studies have shown that it is possible to measure the human antigen-specific T cell response over time through systematic deuterium labeling, but further basic and clinical trials are still needed to determine the optimal timing of surgery (60). Although the results of neoadjuvant immunotherapy on early-stage NSCLC are all from phase I/II clinical trials, they still have reference value. CheckMate 159, LCMC3, and TOP1501 included two cycles of neoadjuvant single agent immunotherapy, and surgery was performed 28 to 56 days after the first cycle, i.e., 1–5 weeks after the end of immunotherapy (26–28). Neoadjuvant immunotherapy combined with chemotherapy (NADIM, NCT02716038) or the combination of two immunotherapies (NEOSTAR) was performed for three to four cycles, and surgery was performed 3–7 weeks after the end of the neoadjuvant immunotherapy (43). In summary, surgery is recommended at 4–6 weeks after the last neoadjuvant immunotherapy cycle.

#### **Consensus 6: There is no definitive evidence that neoadjuvant immunotherapy affects the conduct or safety of surgery**

The morbidity and safety of surgery after neoadjuvant

immunotherapy is still not definitively known. Some clinical trials have demonstrated that after neoadjuvant radiotherapy (NRT) or NCT, surgery is safe and feasible, but that neoadjuvant immunotherapy may lead to tissue adhesion, thus increasing the operative difficulty. Thus, appropriate concerns have been raised as to whether neoadjuvant immunotherapy may increase the surgery complexity (61,62). Currently, the phase I/II studies of neoadjuvant immunotherapy showed that the incidence of any adverse events caused by neoadjuvant single agent immunotherapy was approximately 57%, the incidence of adverse events  $\geq$  grade 3 was approximately 4.5–8%, and the completion rate of planned surgery was 78–100%, which are all similar to those of NCT and NRT (26–28). The NEOSTAR study assessed surgical difficulty and pulmonary function after neoadjuvant immunotherapy, and the results showed that ICIs had less impact on the surgical resection rate and surgical complexity and had no adverse effect on perioperative outcomes (43). A retrospective analysis of 19 patients in the United States showed that, for metastatic or unresectable patients, pneumonectomy after neoadjuvant immunotherapy was feasible, with a high R0 rate, while surgery may be challenging, serious complications were rare (63). There are still no clear conclusions about the impact and safety of neoadjuvant immunotherapy on surgical procedures, and more data are needed.

In summary, based on the available data, there is no conclusive evidence that neoadjuvant immunotherapy adversely affects surgical procedures or their safety.

#### **Consensus 7: pathological remission (MPR, pCR) needs to be assessed, recorded, and reported by specialized pathologist after neoadjuvant immunotherapy**

In a prospective multicenter phase II study of stage IIIA and IIIB NSCLC in 1997, Junker *et al.* (64) developed a tumor regression grading system based on resection specimens (62 cases) from 28 patients with lung squamous-cell carcinoma and 12 patients with lung adenocarcinoma after combined NCT and NRT. The results were compared to spontaneous regressive changes in a control group of 50 untreated NSCLC patients. The proposed three-level grading system of pathological response included grade I (no or only slight tumor regression), grade IIA (significant but incomplete tumor regression, more than 10% vital tumor

tissue), grade IIB (less than 10% vital tumor tissue), and grade III (complete tumor regression without vital tumor tissue). Patients with grade IIB or III tumor regression had a much longer survival time than those with grade II or IIA tumor regression (64). Later, Junker *et al.* improved the grading criteria for tumor regression and defined grade I as no tumor regression or only spontaneous tumor regression, grade II as treatment-induced tumor regression, grade IIA as more than 10% vital tumor tissue, grade IIB as less than 10% vital tumor tissue, and grade III as complete tumor regression. The authors also found that along with the complete resection of the tumor, treatment-induced tumor regression and less than 10% vital tumor tissue were essential for improving long-term outcomes (65). However, many later trials questioned whether the percentage of vital tumor tissue remaining after NCT in NSCLC patients was associated with prognosis. The response of lung squamous-cell carcinoma to NCT was significantly better than that of lung adenocarcinoma, the median percentages of vital tumor tissue were 40% and 60%, respectively, and the critical MPR values were 26% and 12%, respectively (66). In 2017, the College of American Pathologists still recommended MPR as the study endpoint of clinical trials on neoadjuvant immunotherapy for lung cancer. Currently, MPR is defined as neoadjuvant therapy-induced tumor regression with less than 10% vital tumor tissue, and pCR is defined as neoadjuvant therapy-induced complete tumor regression without vital tumor tissue (67).

Because ICIs have only been used in neoadjuvant therapy for NSCLC in recent years, most of the current results come from phase I/II clinical trials. The MPR of neoadjuvant single agent immunotherapy is in the range of 19–45%, and that of immunotherapy combined with neoadjuvant therapy fluctuates within 33–83% (26–28,43). These studies are small, the results have not been verified in phase III clinical trials, and the relationships of MPR with PFS and OS need further confirmation. However, immunotherapy has brought higher MPR and surgical resection rates. The current clinical trials of neoadjuvant immunotherapy can collect more specimens for further exploration.

In summary, pathological remission (MPR, pCR) needs to be assessed, recorded and reported by specialized pathologist after neoadjuvant immunotherapy.

### **Consensus 8: for neoadjuvant immunotherapy in nonprogressive patients, immunotherapy can be resumed after surgery, and it can be maintained for 1 year**

PD-1/PD-L1 inhibitors have greatly altered the treatment strategies of advanced NSCLC. The Chinese Society of Clinical Oncology (CSCO) guidelines recommend that advanced NSCLC patients with positive PD-L1 expression and negative driver mutations can be treated with pembrolizumab monotherapy, and the immunotherapy can be withdrawn at disease progression or 35 cycles (2 years). Advanced NSCLC patients whose tumors do not harbor driver mutations, regardless of PD-L1 expression, can be treated with pembrolizumab monotherapy combined with chemotherapy for 4 cycles, followed by maintenance pembrolizumab monotherapy for a total of 2 years. The CSCO guidelines also recommend 1-year maintenance therapy of durvalumab after concurrent chemoradiotherapy for stage III unresectable NSCLC patients. Currently, there is no definitive recommendation on immune maintenance therapy after neoadjuvant immunotherapy, and it varies from no postoperative immune maintenance therapy to 1-year immune maintenance therapy (26–28,43).

Recently, Antonia *et al.* reported the results of the phase III PACIFIC study (NCT02125461) (68). In this study the role of immune checkpoint blockade with durvalumab in locally advanced, unresectable, stage III NSCLC was evaluated. Eligible patients had NSCLCs without progression after they had been treated with at least two cycles of platinum-based chemotherapy concurrent with radiotherapy (chemo-radiotherapy) at a dose of 54 to 66 Gy. A total of 713 patients were randomly assigned in a 2:1 ratio to receive either durvalumab (10 mg/kg) or placebo every 2 weeks for up to 12 months. Results showed that the co-primary end point of median PFS was 16.8 months in the durvalumab group versus 5.6 months in the placebo group (HR =0.52; 95% CI, 0.42–0.65). In addition, the ORR (assessed by blinded independent central review) was found to be higher in the durvalumab group than in the placebo group (28.4% *vs.* 16.0%,  $P < 0.001$ ). Interestingly, clinical benefit was observed irrespectively of NSCLC tumor stage (IIIA or IIIB), histologic type, or geographic distribution. Most notably, however, brain metastases developed far more frequently in the placebo group as in the durvalumab group

(11.0% vs. 5.5%) (16,17). The OS was also significantly increased with not reached versus 29.1 months (HR =0.68).

Given the experience with immunotherapy in advanced NSCLC and stage III unresectable NSCLC, the neoadjuvant immunotherapy of nonprogressive patients can be resumed after surgery, and can be maintained for 1 year.

**Consensus 9: immunotherapy or induction chemotherapy can be offered in borderline resectable locally advanced NSCLC, and consideration for surgery should be reevaluated upon restaging**

The standard treatment for stage IIIA/B unresectable NSCLC is maintenance therapy with durvalumab monotherapy after concurrent chemoradiotherapy (based on the results of the PACIFIC Trial). Debate exists as to whether these patients are best served by surgery after induction therapy. The ESPATUE study (N =246 of 500 planned, trial had been stopped due to slow recruitment) showed that some patients with stage III unresectable disease may benefit from induction chemotherapy or chemoradiotherapy: their T and N stages were significantly downgraded, and tumors became surgically resectable with a downgrade rate of 44%. Patient characteristics were balanced between the two arms, in which 81 were assigned to surgery (arm A), and 80 were assigned to a chemoradiotherapy boost. In arm B, 81% underwent R0 resection. With a median follow-up after random assignment of 78 months, 5-year OS and PFS did not differ between arms. OS rates of 44% for arm B and 40% for arm A (log-rank P=0.34) and PFS rates of 32% for arm B and 35% for arm A (log-rank P=0.75). OS at 5 years was 34.1% (95% CI, 27.6–40.8%) in all 246 patients, and 216 patients (87.8%) received definitive local treatment (69).

Most recently Reck *et al.* (70) provided the first evidence that the combination of immunotherapy followed by chemotherapy and subsequently maintenance therapy resulted in a significantly improved mOS (CheckMate-9LA). In this phase III trial patients (N=719) were randomized between immunotherapy with ipilimumab and nivolumab (plus two cycles of platinum-based chemotherapy) followed by ipilimumab plus nivolumab as maintenance until progression versus four cycles of platinum-based chemotherapy followed by maintenance therapy (until progression). The ORR rate for the immunotherapy arm was 38% (2% complete remission), mOS rates after one

year were 63% versus 47% (HR =0.66) (approved by FDA and EMA). The results were seen regardless of the PD-L1 status. From this trial the authors concluded that the IO-chemotherapy combination should be considered an effective and well-tolerated novel treatment opportunity for advanced and/or metastatic NSCLC. Although this trial was conducted in stage IV NSCLC patients, the ORR of 38% (including 2% CR rate) is remarkable and further studies are currently planned to evaluate this regimen in earlier stages of NSCLC in the neoadjuvant setting.

Although surgical resection did not increase the postoperative PFS and OS compared to radical chemoradiotherapy, selected patients (T3N2, T4N0/T4N1) had a significant long-term survival benefit in the subgroup analysis, and the benefit in the cohort of patients with stage III B (T4N0-1) patients was the most significant (69). Although many immunotherapy neoadjuvant clinical trials are still underway, the NADIM study indicated that after stage III (N2 or T4N0/N1) patients received four cycles of nivolumab monotherapy combined with chemotherapy, mPFS was found to be 77.1% after 24 months. 30% of patients had treatment-related adverse events of grade 3 or worse; however, none of the adverse events were associated with surgery delays or deaths. The downstage rate was 90.2%, the completion rate of planned surgery was 89.1%, and the MPR rate was 83%, indicating a promising long-term survival benefit in this group of patients (71).

In summary, induction immunotherapy with or without chemotherapy can be considered for unresectable locally advanced NSCLC (particularly in selected N2 and T4 patients), and after the tumor stage is downgraded, the possibility of surgery could be reevaluated. The results of randomized phase III trials with OS as the primary endpoint should be awaited before this strategy is introduced into standard of care.

**Key questions and perspectives**

*Do you use neoadjuvant PD-1/PD-L1 blockade in routine practice? Why or why not?*

**Rafael Rosell:** Neoadjuvant PD-L1 approach is still not standardized, therefore I prefer to include patients in available clinical trials on immune checkpoint inhibitors in combination with chemotherapy. However, understanding that there are patients who may not have access to clinical trials, the use of preoperative neoadjuvant immunotherapy could be adequate for stage IIIA NSCLC patients.

**Mariano Provencio:** I believe that the results of chemo-immunotherapy could support its use in clinical practice, it makes all the same sense as in advanced stages and the results are much better than just chemotherapy. Perhaps we should wait for the ongoing trials, but their adoption should be quick and perhaps accept surrogates of pathological response... but it is not approved in my country.

**Erminia Massarelli:** Yes, I use it within clinical trials of neoadjuvant immunotherapy.

I truly believe that checkpoint inhibitors alone or plus chemotherapy can significantly downstage tumors and I believe the advantage especially in stage II–III NSCLC is significant. However, so far we are only using it within clinical trials.

**Mara B. Antonoff:** We try to enroll as many patients as possible onto trials evaluating neoadjuvant PD-L1 blockade; however, outside of a clinical trial, we do not routinely use it in the neoadjuvant setting as it is not yet approved for use outside of a trial.

**Toyoaki Hida:** We don't use neoadjuvant PD-1/PD-L1 blockade in routine practice, because immune checkpoint inhibitors (ICIs) are not yet approved for the use of preoperative treatment in lung cancer in Japan. We use ICIs for preoperative treatment in clinical trials only.

**Marc de Perrot:** I do not use neoadjuvant PD-L1 blockade in routine practice due to the lack of access. Neoadjuvant IO is part of clinical trials in our institution.

**Steven H. Lin:** No, all our patients with this approach are treated on clinical trial, since it is currently not a standard approach for the use of neoadjuvant immunotherapy although early studies have shown promising major pathologic response. All neoadjuvant approaches should be done on the ongoing numerous single arm or randomized trials. Neoadjuvant chemotherapy is still a good option as the control arm in those randomized trials.

**Massimo Di Maio:** We do not currently use neoadjuvant PD-1/PD-L1 blockade in routine practice, because this is not a standard approach according to our national guidelines by AIOM (Italian Association of Medical Oncology). In general, neoadjuvant treatment is not commonly used for patients who are judged eligible for surgery, with the exception of a known positivity of mediastinal lymph nodes, when a neoadjuvant chemotherapy is commonly considered before re-evaluation with the surgeon.

As a general rule, Italy has a publicly funded health system, and only treatments approved and reimbursed can be considered in clinical practice. Furthermore, AIOM has chosen to produce recommendations only for

drugs/ treatments that are reimbursed by the national health system, considering that it would make no sense to recommend something that cannot be used in clinical practice.

At the moment, use of neoadjuvant PD-1/PD-L1 blockade is limited to participation in clinical trials.

**Antonio Rossi:** I do not use neoadjuvant PD-1/PD-L1 blockade in routine practice because, despite the very interesting results coming from phase I/II trials, it is still an experimental approach. The Site in which I worked had no possibility to be involved in trials addressing this issue in order to participate to the investigation of this strategic approach. Randomized trials are ongoing and if the results will be positive this approach should be evaluated by regulatory agencies and then considered for clinical practice.

**Dirk de Ruyscher:** No, only in the context of a clinical trial. There are no phase III data showing an increase of OS.

**Robert A. Ramirez:** No. The data looks promising for combination chemo/IO in the neoadjuvant setting, however, outside of a clinical trial we don't have large scale studies to guide us. Plus, in the US we run into reimbursement issues and cannot get this in the neoadjuvant setting.

**Wolfram C. M. Dempke:** In our institution neoadjuvant therapies are currently not used routinely as they are not FDA/EMA approved so far (with very few exceptions). However, we participate in clinical studies.

**D. Ross Camidge:** NO—It is not licensed and its OS benefit is not clear. It needs to be compared with adjuvant therapy as it will cause delay in surgery.

**Nicolas Guibert:** I do, but only within clinical trials because no ICIs as neoadjuvant therapies are approved in France yet.

**Raffaele Califano:** It's not in routine practice as not approved/funded.

*Do you think biomarker is necessary in choosing neoadjuvant PD-1/PD-L1 blockade therapy? If yes, which ones?*

**Rafael Rosell:** Needless to say, immune biomarkers are desirable and a PD-L1 tumor proportion score (TPS) of more than 50% could serve and encourage patients. However, patients with a TPS of lower than 50%, but more than 1%, are also suitable for the neoadjuvant immunotherapy approach. Furthermore, if liquid biopsy is accessible, a high blood TMB is also orientative to predict

response to immune checkpoint inhibitors. NSCLC with driver mutations or gene fusions cannot be considered for neoadjuvant immunotherapy approaches.

**Mariano Provencio:** No, I don't think so.

**Erminia Massarelli:** No, I do not think current biomarkers are needed considering that PD-L1 negative patient have good benefit as well. I believe we still need to discover reliable markers of response in PD-L1 $\leq$ 50%.

**Mara B. Antonoff:** We know that PD-L1 status can help determine patients who are likely to do well, and we also know that patients with EGFR and ALK mutations are not likely to derive benefit, and would not include such patients on these trials when they can receive targeted therapy instead.

**Toyoaki Hida:** Yes: We think biomarker may be necessary in choosing neoadjuvant PD-1/PD-L1 blockade therapy in the case of single ICI use.

Candidate for biomarker: PD-1 CD8 T cells, PD-1 regulatory T (Treg) cells, frequency of PD-1 CD8 T cells relative to that of PD-1 Treg cells.

It may be not necessary in the cases of ICI use in combination with platinum-doublet chemotherapy, although it would be better to have biomarkers that can predict the effect.

**Marc de Perrot:** PD-L1 expression on tumor cell is a good biomarker when PD-1/PD-L1 is used as single therapy. However, biomarker may not be necessary when PD-1/PD-L1 blockade is combined with chemotherapy.

**Steven H. Lin:** We select for patients who shouldn't get immunotherapy, which are patients who have an oncogene driven mutation such as EGFR, ALK, ROS, RET. We don't select for other markers like PD-L1 status.

**Massimo Di Maio:** Very good question. It probably depends if we are talking about single agent PD-1/PD-L1 blockade or combination with chemotherapy. If combined with chemotherapy, based on the results obtained in the advanced setting, we could avoid a pre-treatment selection according to PD-L1 expression. In fact, we know that chemotherapy is an acceptable and effective strategy as neoadjuvant treatment, so the risk of undertreatment would be limited. On the other hand, in the case of single-agents, we know that a not negligible proportion of patients would be at risk of not obtaining a response, so a selection should be probably considered (with the only exception of a very short planned duration of treatment, which would not delay the timing of surgery), probably based on the PD-L1 expression. Unfortunately, evidence about biomarkers' role in this setting (including PD-L1 expression but also TMB)

is not robust.

**Antonio Rossi:** Biomarkers should drive the treatment for any stage of NSCLC disease. Unfortunately, we do not have biomarkers for every subgroup of NSCLC patients. However, with the available data, in the neoadjuvant setting, no specific biomarker can help in defining the subgroup of patients which could much benefit by neoadjuvant immunotherapy. The potential role of PD-L1 expression and other potential biomarkers such as MSI or TMB, in this setting should be evaluated more extensively in the ongoing randomized trials, in order to have the possibility to select NSCLC patients who could much benefit from neoadjuvant immunotherapy.

**Dirk de Ruyscher:** At present, we do not have a good marker for concurrent chemo-PD-1/PD-L1 blockade, while for anti-PD-1/PD-L1 alone, tumor PD-L1 expression is still the best. There are many new developments such as the T-cell (CD8 and Treg) density on the initial tumor biopsy, which are promising.

**Robert A. Ramirez:** No. I suppose if someone were PD-L1 >50%, MSI high or high TMB that may sway me to use single agent IO, however, we run into lack of data as above. We also know the RR in the combination chemo/IO is higher in the metastatic setting regardless of biomarkers.

**Wolfram C. M. Dempke:** This is a controversial discussion. We follow the label of the IOs (have added this in the manuscript as well), but truly believe that additional biomarkers are urgently warranted (for new examples see my paragraph within the manuscript).

**D. Ross Camidge:** I suspect it may influence outcomes – the obvious cases would be ALK and EGFR.

**Nicolas Guibert:** No clear biomarker of response (including TMB and PD-L1) has been identified yet in the neoadjuvant setting like it has been well stated in the manuscript.

I would thus not recommend the use of biomarker to guide neoadjuvant ICI yet.

I would, however, like proposed in the manuscript, suggest to avoid neoadjuvant ICI monotherapy in patients with EGFR, ALK or STK11 alterations, given the low response rates observed in these populations in advanced stages.

This extrapolation may however need to be validated in clinical trials.

More endeavors are however needed to better select patients.

**Raffaele Califano:** PD-L1 using single agent PD-1/PD-L1 blockade. No need for biomarker if using neoadjuvant

Chemo + IO.

*How do/will you determine the cycles of neoadjuvant PD-1/PD-L1 blockade therapy?*

**Rafael Rosell:** At present, there is no pre-specified number of cycles. Two to four cycles are the recommendation.

**Mariano Provencio:** We use 3 cycles.

**Erminia Massarelli:** Usually Two to three cycles as per clinical trials guidelines.

**Mara B. Antonoff:** We believe anywhere from 2–4 cycles is beneficial, and the data don't yet define an optimal number. We don't use it outside of a clinical trial, so at this time, we use it for the designated amount of time based on the clinical trial. We need further data from these studies to decide the exact number of cycles.

**Toyoaki Hida:** 2-3 cycles depending on the degree of bone marrow suppression (degree of irAE).

**Marc de Perrot:** Two to four cycles are sufficient in the neoadjuvant setting.

**Steven H. Lin:** Standardly it has been as little as one cycle to 2 cycles, +/- chemotherapy. Neoadjuvant chemoradiation with IO is being explored as well.

**Massimo Di Maio:** I would determine the cycles of neoadjuvant PD-1/PD-L1 blockade according to the evidence produced by clinical trials.

In the case of combination with chemotherapy, I would have no problem to plan 3 or 4 cycles of treatment, considering that the risk of undertreatment should be negligible. In the case of immunotherapy alone, it would be necessary to avoid a long treatment that could increase the risk of disease progression. In this latter case, as a rule of thumb, I would limit the treatment to 1 month/1.5 months.

**Antonio Rossi:** Based on the available data and the design of the ongoing randomized trials, 2–4 cycles of neoadjuvant single-agent immunotherapy or combined with platinum-based chemotherapy should be appropriate.

**Dirk de Ruyscher:** The best for the future may be ctDNA, but this needs more development and especially standardization. Advanced quantitative imaging methods based on e.g., deep learning algorithms show great promise.

**Robert A. Ramirez:** I agree that 2 cycles of neoadjuvant treatment be followed by imaging and a multidisciplinary review to see when the right time for surgery should be. If responding following 2 cycles but still borderline resectable, then an additional 2 cycles could be beneficial.

**Wolfram C. M. Dempke:** It depends on the early responses seen—if there is a response, we normally go for 4

cycles with a re-evaluation after two cycles.

**D. Ross Camidge:** To date 2–4 cycles are given but the stress of delaying surgery has to be weighed against data.

**Nicolas Guibert:** The number of cycles will depend on the approvals obtained in France. It will follow the regimen studied in the RCT that will lead to this approval. The most appealing data have however been observed with ICI and chemo combinations, with 3 to 4 cycles.

**Raffaele Califano:** We will consider a cycle 3 or 4 weeks of treatment.

*What adjuvant treatment regimen do/will you choose among those respond to neoadjuvant PD-1/PD-L1 blockade therapy?*

**Rafael Rosell:** For responding patients with complete or major pathological response, adjuvant immunotherapy for 1 year could be recommended.

**Mariano Provencio:** We use carbo-taxol-nivolumab.

**Erminia Massarelli:** I continue same immunotherapy plus or minus chemotherapy. However if MPR is significant I only continue immunotherapy if patients were treated with IO alone as induction.

**Mara B. Antonoff:** This depends on the stage of disease. For those with early-stage disease, no need for adjuvant therapy. However, for recurrent disease, node positive disease, or large tumors that would otherwise get adjuvant systemic therapy, immunotherapy should be given to those with elevated PD-L1 expression.

**Toyoaki Hida:** We choose ICI used in the preoperative treatment.

**Marc de Perrot:** PD-1/PD-L1 blockade will be maintained in the adjuvant setting.

**Steven H. Lin:** Although chemotherapy is not used adjuvant even if it is given concurrently with immunotherapy, only the immunotherapy as a single agent is used in the adjuvant setting for most trials. Radiotherapy is used per the discretion of the trial, with some global trials excluding the use of postoperative radiotherapy for IIIA-N2 disease resected to negative margins.

**Massimo Di Maio:** We have no data to produce evidence-based recommendations. For patients who have not received chemotherapy, I would consider adjuvant chemotherapy according to current guidelines. For patients who have received chemotherapy as neoadjuvant treatment (i.e., combo of chemo + immuno), probably they could receive no further systemic treatment after surgery. I prefer waiting for clinical trials.

**Antonio Rossi:** Platinum-based doublets, if not received in neoadjuvant setting, should be considered the standard-of-care with or without PD-1/PD-L1 inhibitors as adjuvant treatment. At the moment, this combination approach is reserved for patients enrolled in clinical trials.

**Dirk de Ruyscher:** In case of an MPR or a pCR: the same anti-PD-1/PD-L1 as pre-operatively. In case no MPR or pCR was achieved: (I) if only neoadjuvant immunotherapy: 4 cycles of a platinum doublet depending on the histology of the tumor; (II) if concurrent chemotherapy and immunotherapy: no further adjuvant therapy or no systemic therapy has shown a benefit.

**Robert A. Ramirez:** If the patient has not received neoadjuvant chemo and only received IO then if adjuvant therapy is indicated then I would use a cisplatin-based regimen based on histology.

**Wolfram C. M. Dempke:** We believe that the adjuvant treatment regimen should be the same as used for the neoadjuvant setting, however without chemotherapy if already used in the neoadjuvant part.

**D. Ross Camidge:** Trial specific at present.

**Nicolas Guibert:** Again, the adjuvant regimen will depend on the approvals obtained in France that will follow the main RCTs' outcomes. Most trials were however designed with ana approximately 1-year adjuvant immunotherapy (13 additional pembrolizumab adjuvant cycles in the KEYNOTE 617, 16 cycles of atezolizumab in the IMPOWER 030) and it is very likely that this approach may be the most suited. It will also certainly depend on the pathological response (CR, MPR) and definitive pathological stage (pTNM) after surgery.

**Raffaele Califano:** We will give platinum/vinorelbine ×4 cycle.

## Acknowledgments

The authors appreciate the academic support from AME Lung Cancer Collaborative Group.

*Funding:* None.

## Footnote

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tlcr-2020-63>). The authors have no conflicts of interest to declare. The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424.
2. Chen W, Zheng R, Baade PD, et al. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016;66:115-32.
3. Molina JR, Yang P, Cassivi SD, et al. Non-small cell lung cancer: epidemiology, risk factors, treatment, and survivorship. *Mayo Clin Proc* 2008;83:584-94.
4. Arriagada R, Bergman B, Dunant A, et al. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *N Engl J Med* 2004;350:351-60.
5. Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. *N Engl J Med* 2005;352:2589-97.
6. Douillard JY, Rosell R, De Lena M, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB-IIIa non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. *Lancet Oncol* 2006;7:719-27.
7. Goldstraw P, Chansky K, Crowley J, et al. The IASLC Lung Cancer Staging Project: Proposals for Revision of the TNM Stage Groupings in the Forthcoming (Eighth) Edition of the TNM Classification for Lung Cancer. *J Thorac Oncol* 2016;11:39-51.
8. Uprety D, Mandrekar SJ, Wigle D, et al. Neoadjuvant Immunotherapy for NSCLC: Current Concepts and Future Approaches. *J Thorac Oncol* 2020;15:1281-97.

9. NSCLC Meta-analysis Collaborative Group. Preoperative chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual participant data. *Lancet* 2014;383:1561-71.
10. Lim E, Harris G, Patel A, et al. Preoperative versus postoperative chemotherapy in patients with resectable non-small cell lung cancer: systematic review and indirect comparison meta-analysis of randomized trials. *J Thorac Oncol* 2009;4:1380-8.
11. Gandhi L, Rodriguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;378:2078-92.
12. Paz-Ares L, Luft A, Vicente D, et al. Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;379:2040-51.
13. Mok TSK, Wu YL, Kudaba I, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819-30.
14. Hanna NH, Schneider BJ, Temin S, et al. Therapy for Stage IV Non-Small-Cell Lung Cancer Without Driver Alterations: ASCO and OH (CCO) Joint Guideline Update. *J Clin Oncol* 2020;38:1608-32.
15. Santarpia M, Aguilar A, Chaib I, et al. Non-Small-Cell Lung Cancer Signaling Pathways, Metabolism, and PD-1/PD-L1 Antibodies. *Cancers (Basel)* 2020;12:1475.
16. Garon EB, Hellmann MD, Rizvi NA, et al. Five-Year Overall Survival for Patients With Advanced Non-Small-Cell Lung Cancer Treated With Pembrolizumab: Results From the Phase I KEYNOTE-001 Study. *J Clin Oncol* 2019;37:2518-27.
17. Topalian SL, Hodi FS, Brahmer JR, et al. Five-Year Survival and Correlates Among Patients With Advanced Melanoma, Renal Cell Carcinoma, or Non-Small Cell Lung Cancer Treated With Nivolumab. *JAMA Oncol* 2019;5:1411-20.
18. O'Donnell JS, Hoefsmit EP, Smyth MJ, et al. The Promise of Neoadjuvant Immunotherapy and Surgery for Cancer Treatment. *Clin Cancer Res* 2019;25:5743-51.
19. Roth JA, Fossella F, Komaki R, et al. A randomized trial comparing perioperative chemotherapy and surgery with surgery alone in resectable stage IIIA non-small-cell lung cancer. *J Natl Cancer Inst* 1994;86:673-80.
20. Rosell R, Gomez-Codina J, Camps C, et al. A randomized trial comparing preoperative chemotherapy plus surgery with surgery alone in patients with non-small-cell lung cancer. *N Engl J Med* 1994;330:153-8.
21. Song WA, Zhou NK, Wang W, et al. Survival benefit of neoadjuvant chemotherapy in non-small cell lung cancer: an updated meta-analysis of 13 randomized control trials. *J Thorac Oncol* 2010;5:510-6.
22. Keir ME, Butte MJ, Freeman GJ, et al. PD-1 and its ligands in tolerance and immunity. *Annu Rev Immunol* 2008;26:677-704.
23. Liu J, Blake SJ, Yong MC, et al. Improved Efficacy of Neoadjuvant Compared to Adjuvant Immunotherapy to Eradicate Metastatic Disease. *Cancer Discov* 2016;6:1382-99.
24. Bourgeois-Daigneault MC, Roy DG, Aitken AS, et al. Neoadjuvant oncolytic virotherapy before surgery sensitizes triple-negative breast cancer to immune checkpoint therapy. *Sci Transl Med* 2018;10:eaao1641.
25. Brooks J, Fleischmann-Mundt B, Woller N, et al. Perioperative, Spatiotemporally Coordinated Activation of T and NK Cells Prevents Recurrence of Pancreatic Cancer. *Cancer Res* 2018;78:475-88.
26. Forde PM, Chaft JE, Smith KN, et al. Neoadjuvant PD-1 Blockade in Resectable Lung Cancer. *N Engl J Med* 2018;378:1976-86.
27. Kwiatkowski DJ RV, Chaft JE, BE J. Neoadjuvant atezolizumab in resectable nonsmall cell lung cancer (NSCLC): Interim analysis and biomarker data from a multicenter study (LCMC3). *J Clin Oncol* 2019;37:abstr 8503.
28. Ready N, Tong B, Clarke J, et al. Neoadjuvant Pembrolizumab in Early Stage Non-Small Cell Lung Cancer (NSCLC): Toxicity, Efficacy, and Surgical Outcomes. *J Thorac Oncol* 2019;14:S745.
29. Wislez M, Mazieres J, Lavole A, et al. Neoadjuvant durvalumab in resectable non-small cell lung cancer (NSCLC): preliminary results from a multicenter study (IFCT-1601 IONESCO). *Ann Oncol* 2020;31:S794.
30. Gao S, Li N, Gao S, et al. Neoadjuvant PD-1 inhibitor (Sintilimab) in NSCLC. *J Thorac Oncol* 2020;15:816-26.
31. Besse B, Adam J, Cozic N, et al. Neoadjuvant atezolizumab (A) for resectable non-small cell lung cancer (NSCLC): results from the phase II PRINCEPS trial. *Ann Oncol* 2020;31:S735-43.
32. Shu CA, Gainor JF, Awad MM, et al. Neoadjuvant atezolizumab and chemotherapy in patients with resectable non-small-cell lung cancer: an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2020;21:786-95.
33. Provencio M, Nadal E, Insa A, et al. Neoadjuvant chemotherapy and nivolumab in resectable non-small-cell

- lung cancer (NADIM): an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2020;21:1413-22.
34. Xia L, Liu Y, Wang Y. PD-1/PD-L1 Blockade Therapy in Advanced Non-Small-Cell Lung Cancer: Current Status and Future Directions. *Oncologist* 2019;24:S31-S41.
  35. Reck M, Rodriguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2016;375:1823-33.
  36. Spigel D, de Marinis F, Giaccone G, et al. IMpower110: Interim overall survival (OS) analysis of a phase III study of atezolizumab (atezo) vs platinum-based chemotherapy (chemo) as first-line (1L) treatment (tx) in PD-L1-selected NSCLC. *Ann Oncol* 2019;30:v851-v934.
  37. Available online: <https://wwwfdagov/news-events/press-announcements/fda-approves-first-cancer-treatment-any-solid-tumor-specific-genetic-feature>
  38. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor Mutational Burden. *N Engl J Med* 2018;378:2093-104.
  39. Available online: <https://wwwfdagov/drugs/drug-approvals-and-databases/fda-approves-pembrolizumab-adults-and-children-tmb-h-solid-tumors>
  40. Cappuzzo F, Reck M, Socinski MA, et al. IMpower150: Exploratory analysis of brain metastases development. *J Clin Oncol* 2020;38:abstr 9587.
  41. Zhou J, Tang Z, Gao S, et al. Tumor-Associated Macrophages: Recent Insights and Therapies. *Front Oncol* 2020;10:188.
  42. Ford K, Hanley CJ, Mellone M, et al. NOX4 Inhibition Potentiates Immunotherapy by Overcoming Cancer-Associated Fibroblast-Mediated CD8 T-cell Exclusion from Tumors. *Cancer Res* 2020;80:1846-60.
  43. Tina Cascone, William Nassib William, Annikka Weissferdt, et al. Neoadjuvant nivolumab (N) or nivolumab plus ipilimumab (NI) for resectable non-small cell lung cancer (NSCLC): Clinical and correlative results from the NEOSTAR study. *J Clin Oncol* 2019;37:abstr 8504.
  44. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144:646-74.
  45. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med* 2015;373:1627-39.
  46. Cascone T, William WN, Weissferdt A, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 2016;387:1540-50.
  47. Bylicki O, Paleiron N, Margery J, et al. Targeting the PD-1/PD-L1 Immune Checkpoint in EGFR-Mutated or ALK-Translocated Non-Small-Cell Lung Cancer. *Target Oncol* 2017;12:563-9.
  48. Skoulidis F, Arbour KC, Hellmann MD, et al. Association of STK11/LKB1 genomic alterations with lack of benefit from the addition of pembrolizumab to platinum doublet chemotherapy in non-squamous non-small cell lung cancer. *J Clin Oncol* 2019;37:abstr 102.
  49. Mutations Associated with Sensitivity or Resistance to Immunotherapy in mNSCLC. Analysis from the MYSTIC Trial. Available online: [https://www.eurekalert.org/pub\\_releases/2019-09/iaft-maw090819.php](https://www.eurekalert.org/pub_releases/2019-09/iaft-maw090819.php)
  50. Cho BC, Lopes G, Kowalski DM, et al. Relationship between STK11 and KEAP1 mutational status and efficacy in KEYNOTE-042: pembrolizumab monotherapy versus platinum-based chemotherapy as first-line therapy for PD-L1-positive advanced NSCLC. *AACR* 2020;80:abstr CT084.
  51. Nadal E, Heeke S, Benzaquen J, et al. Two Patients With Advanced-Stage Lung Adenocarcinoma With Radiologic Complete Response to Nivolumab Treatment Harboring an STK11/LKB1 Mutation. *JCO Precis Oncol* 2020;4:1239-45.
  52. Uprety D, Mandrekar SJ, Wigle D, et al. Neoadjuvant Immunotherapy for NSCLC: Current Concepts and Future Approaches. *J Thorac Oncol* 2020;15:1281-97.
  53. Liu J, O'Donnell JS, Yan J, et al. Timing of neoadjuvant immunotherapy in relation to surgery is crucial for outcome. *Oncoimmunology* 2019;8:e1581530.
  54. Amaria RN, Menzies AM, Burton EM, et al. Neoadjuvant systemic therapy in melanoma: recommendations of the International Neoadjuvant Melanoma Consortium. *Lancet Oncol* 2019;20:e378-89.
  55. Travis WD, Dacic S, Wistuba I, et al. IASLC Multidisciplinary Recommendations for Pathologic Assessment of Lung Cancer Resection Specimens After Neoadjuvant Therapy. *J Thorac Oncol* 2020;15:709-40.
  56. William WN Jr, Pataer A, Kalhor N, et al. Computed tomography RECIST assessment of histopathologic response and prediction of survival in patients with resectable non-small-cell lung cancer after neoadjuvant chemotherapy. *J Thorac Oncol* 2013;8:222-8.
  57. Tao X, Li N, Wu N, et al. The efficiency of (18)F-FDG PET-CT for predicting the major pathologic response to the neoadjuvant PD-1 blockade in resectable non-

- small cell lung cancer. *Eur J Nucl Med Mol Imaging* 2020;47:1209-19.
58. Huang C, Liu S, Tong X, et al. Extracellular vesicles and ctDNA in lung cancer: biomarker sources and therapeutic applications. *Cancer Chemother Pharmacol* 2018;82:171-83.
  59. Chae YK, Oh MS. Detection of Minimal Residual Disease Using ctDNA in Lung Cancer: Current Evidence and Future Directions. *J Thorac Oncol* 2019;14:16-24.
  60. Akondy RS, Fitch M, Edupuganti S, et al. Origin and differentiation of human memory CD8 T cells after vaccination. *Nature* 2017;552:362-7.
  61. Edelman MJ, Suntharalingam M, Burrows W, et al. Phase I/II trial of hyperfractionated radiation and chemotherapy followed by surgery in stage III lung cancer. *Ann Thorac Surg* 2008;86:903-10.
  62. Shah AA, Berry MF, Tzao C, et al. Induction chemoradiation is not superior to induction chemotherapy alone in stage IIIA lung cancer. *Ann Thorac Surg* 2012;93:1807-12.
  63. Bott MJ, Cools-Lartigue J, Tan KS, et al. Safety and Feasibility of Lung Resection After Immunotherapy for Metastatic or Unresectable Tumors. *Ann Thorac Surg* 2018;106:178-83.
  64. Junker K, Thomas M, Schulmann K, et al. Tumour regression in non-small-cell lung cancer following neoadjuvant therapy. Histological assessment. *J Cancer Res Clin Oncol* 1997;123:469-77.
  65. Junker K, Langner K, Klinke F, et al. Grading of tumor regression in non-small cell lung cancer : morphology and prognosis. *Chest* 2001;120:1584-91.
  66. Qu Y, Emoto K, Eguchi T, et al. Pathologic Assessment After Neoadjuvant Chemotherapy for NSCLC: Importance and Implications of Distinguishing Adenocarcinoma From Squamous Cell Carcinoma. *J Thorac Oncol* 2019;14:482-93.
  67. Owen D, Chaft JE. Immunotherapy in surgically resectable non-small cell lung cancer. *J Thorac Dis* 2018;10:S404-S11.
  68. Antonia SJ, Villegas A, Daniel D, et al. Durvalumab after Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer. *N Engl J Med* 2017;377:1919-29.
  69. Eberhardt WE, Pottgen C, Gauler TC, et al. Phase III Study of Surgery Versus Definitive Concurrent Chemoradiotherapy Boost in Patients With Resectable Stage IIIA(N2) and Selected IIIB Non-Small-Cell Lung Cancer After Induction Chemotherapy and Concurrent Chemoradiotherapy (ESPA-TUE). *J Clin Oncol* 2015;33:4194-201.
  70. Reck M, Ciuleanu T, Cobo M, et al. First-line nivolumab (NIVO) + ipilimumab (IPI) combined with 2 cycles of platinum-based chemotherapy (chemo) vs 4 cycles of chemo in advanced non-small cell lung cancer (NSCLC): Patient-reported outcomes (PROs) from CheckMate 9LA. *Ann Oncol* 2020;31:S1142-215.
  71. Provencio M, Nadal E, Insa A, et al. NEO-adjuvant chemo-immunotherapy for the treatment of STAGE IIIA resectable non-small-cell lung cancer (NSCLC): A phase II multicenter exploratory study-Final data of patients who underwent surgical assessment. *J Clin Oncol* 2019;37:abstr 8509.

**Cite this article as:** Liang W, Cai K, Chen C, Chen H, Chen Q, Fu J, Hu J, Jiang T, Jiao W, Li S, Liu C, Liu D, Liu W, Liu Y, Ma H, Pan X, Qiao G, Tian H, Wei L, Zhang Y, Zhao S, Zhao X, Zhou C, Zhu Y, Zhong R, Li F, Rosell R, Provencio M, Massarelli E, Antonoff MB, Hida T, de Perrot M, Lin SH, Di Maio M, Rossi A, De Ruyscher D, Ramirez RA, Dempke WCM, Camidge DR, Guibert N, Califano R, Wang Q, Ren S, Zhou C, He J. Expert consensus on neoadjuvant immunotherapy for non-small cell lung cancer. *Transl Lung Cancer Res* 2020;9(6):2696-2715. 10.21037/tlcr-2020-63

Table S1 The ongoing trials of neoadjuvant immunotherapy

Study title	NCT.no	Key inclusion and exclusion criteria	Drug	Control group	Primary and secondary endpoint
A Single-arm, Phase II Study of Neoadjuvant MPDL3280A, Nab-paclitaxel and Carboplatin (MAC) in Resectable Non-small Cell Lung Cancer (NSCLC)	NCT02716038	Eligible patients were aged 18 years or older and had resectable American Joint Committee on Cancer-defined stage IB-IIIa non-small-cell lung cancer, an ECOG performance status of 0-1, and a history of smoking exposure	Atezolizumab; carboplatin; nab-paclitaxel	Carboplatin + nab-paclitaxel	The primary endpoint was major pathological response, defined as the presence of 10% or less residual viable tumour at the time of surgery
Randomized, Open-label, Controlled Phase III Trial Comparing Pembrolizumab-platinum Based Chemotherapy Combination With Pembrolizumab Monotherapy in First Line Treatment of Non-small-cell Lung Cancers (NSCLC) With PDL1 Expression $\geq$ 50%	NCT04547504	Advanced NSCLC molecularly defined by a PDL1 expression $\geq$ 50% of tumour cells and no EGFR mutations or ALK rearrangement	Pembrolizumab; cisplatin; carboplatin AUC	Pembrolizumab alone	The primary endpoint was PFS. The secondary endpoints were PFS, iPFS, ORR, OS, DOT, DOR, AE and AESI
A Phase 3, Randomized, Double-blind Trial of Pembrolizumab (MK-3475) With or Without Lenvatinib (E7080/MK-7902) in Participants With Treatment-naïve, Metastatic Non-small Cell Lung Cancer (NSCLC) Whose Tumors Have a Tumor Proportion Score (TPS) Greater Than or Equal to 1% (LEAP-007)-China Extension Study	NCT04676412	No prior systemic therapy for their metastatic NSCLC whose tumors has a PD-L1 TPS $\geq$ 1%	Lenvatinib; pembrolizumab	Pembrolizumab + placebo	The primary endpoints were PFS and OS. The secondary endpoints were ORR, AE, GHS, QoL, EORTC and TTD
A Phase 3, Randomized, Double-blind Trial of Pembrolizumab (MK-3475) With or Without Lenvatinib (E7080/MK-7902) in Participants With Treatment-naïve, Metastatic Non-small Cell Lung Cancer (NSCLC) Whose Tumors Have a Tumor Proportion Score (TPS) Greater Than or Equal to 1% (LEAP-007)	NCT03829332	No prior systemic therapy for their metastatic NSCLC whose tumors has a PD-L1 TPS $\geq$ 1%	Lenvatinib; pembrolizumab	Pembrolizumab + placebo	The primary endpoints were PFS and OS. The secondary endpoints were ORR, AE, GHS, QoL, EORTC and TTD
A Randomized, Phase 3 Trial With Anti-PD-1 Monoclonal Antibody Pembrolizumab (MK-3475) Versus Placebo for Patients With Early Stage NSCLC After Resection and Completion of Standard Adjuvant Therapy (PEARLS)	NCT02504372	Stage IB/II-IIIa NSCLC who have undergone surgical resection	Pembrolizumab	Placebo	The primary endpoint was DFS. The secondary endpoints were OS and LCSS
A Phase 3, Randomized, Placebo-Controlled Clinical Study to Evaluate the Safety and Efficacy of Stereotactic Body Radiotherapy (SBRT) With or Without Pembrolizumab (MK-3475) in Participants With Medically Inoperable Stages I or IIA Non Small Cell Lung Cancer (NSCLC) (KEYNOTE-867)	NCT03924869	Medically inoperable Stage I or IIA NSCLC	Pembrolizumab; SBRT	SBRT + placebo	The primary endpoints were EFS and OS. The secondary endpoints were TDDM, AE and EORTC QLQ-LC13/-C30 SCORE
A Phase 3, Multicenter, Randomized, Open-label Trial to Compare the Efficacy and Safety of Pembrolizumab (MK-3475) in Combination With Lenvatinib (E7080/MK-7902) Versus Docetaxel in Previously Treated Participants With Metastatic Non-small Cell Lung Cancer (NSCLC) and Progressive Disease (PD) After Platinum Doublet Chemotherapy and Immunotherapy (LEAP-008)	NCT03976375	Metastatic NSCLC and progressive disease (PD) after platinum doublet chemotherapy and treatment with one prior anti-PD-1/PD-L1	Pembrolizumab; lenvatinib; docetaxel	Docetaxel	The primary endpoints were OS and PFS. The secondary endpoints were ORR, DOR, AE, EORTC QLQ-LC13/-C30 SCORE and TTD in EORTC QLQ-LC13 Cough (Item 31) Scale Score
A Randomized, Double-Blind, Phase 3 Study of Pemetrexed + Platinum Chemotherapy With or Without Pembrolizumab (MK-3475) in TKI-resistant EGFR-mutated Tumors in Metastatic Non-squamous Non-small Cell Lung Cancer (NSCLC) Participants (KEYNOTE-789)	NCT03515837	Adults with the following types of TKI-resistant, EGFR-mutated, metastatic NSCLC tumors: (I) TKI-failures (including osimertinib [TAGRISSO <sup>®</sup> ] failure) with T790M-negative mutation tumors, (II) T790M-positive mutation tumors with prior exposure to osimertinib, and (III) first-line osimertinib failure regardless of T790M mutation status	Pembrolizumab; pemetrexed; carboplatin; cisplatin	Placebo + pemetrexed + carboplatin+ cisplatin	The primary endpoints were OS and PFS. The secondary endpoints were ORR, DOR, AE, EORTC QLQ-LC13/-C30 SCORE and TTD in the EORTC Questionnaire Composite Endpoint of Cough, Chest Pain or Dyspnea
An Adaptive Phase III, Multicenter, Randomized, Open-Label, Controlled Study of M7824 (Bintrafusp Alfa) Versus Pembrolizumab as a First-line Treatment in Patients With PD-L1 Expressing Advanced Non-small Cell Lung Cancer	NCT03631706	Advanced NSCLC with high PD-L1-tumor expression, with no EGFR mutation or ALK translocation	M7824; pembrolizumab	Pembrolizumab	The primary endpoints were OS and PFS. The secondary endpoint were TEAEs, NCI-CTCAE, CR, PR and Immunogenicity
A Phase III, Randomized, Double-blind Trial of Platinum Doublet Chemotherapy +/-Pembrolizumab (MK-3475) as Neoadjuvant/Adjuvant Therapy for Participants With Resectable Stage II, IIIA, and Resectable IIIB (T3-4N2) Non-small Cell Lung Cancer (NSCLC) (KEYNOTE-671)	NCT03425643	Resectable stage II, IIIA, and resectable IIIB (T3-4N2) NSCLC	Pembrolizumab; cisplatin; gemcitabine; pemetrexed	NAC + neoadjuvant/adjuvant placebo	The primary endpoints were EFS and OS. The secondary endpoints were mPR rate, pCR rate, AE and EGHs/QoL SCORE
A Randomized, Double-blind, Placebo-controlled, Phase III Study Evaluating the Efficacy and Safety of Pembrolizumab Plus Platinum-based Doublet Chemotherapy With or Without Canakinumab as First Line Therapy for Locally Advanced or Metastatic Non-squamous and Squamous Non-small Cell Lung Cancer Subjects (CANOPY-1)	NCT03631199	Previously untreated locally advanced or metastatic non-squamous and squamous NSCLC subjects	Canakinumab; pembrolizumab; carboplatin; cisplatin; paclitaxel; nab-paclitaxel; pemetrexed	Canakinumab matching-placebo	The primary endpoint were DLTs, PFS and OS. The secondary endpoints were ORR, DCR, DOR and TTR
A Phase 3 Randomized, Placebo-controlled Study to Evaluate the Safety and Efficacy of Pemetrexed + Platinum Chemotherapy + Pembrolizumab (MK-3475) With or Without Lenvatinib (E7080/MK-7902) as First-line Intervention in Participants With Metastatic Nonsquamous Non-small Cell Lung Cancer (LEAP-006)	NCT03829319	Adults with metastatic nonsquamous NSCLC	Pembrolizumab; carboplatin; cisplatin; pemetrexed; lenvatinib	Pemetrexed + platinum chemotherapy + pembrolizumab + placebo	The primary endpoints were DLTs, AES, PFS and OS. The secondary endpoints were ORR, DOR and EORTC QLQ-LC13/-C30 SCORE
LIBRETTO-431: A Multicenter, Randomized, Open-Label, Phase 3 Trial Comparing Selpercatinib to Platinum-Based and Pemetrexed Therapy With or Without Pembrolizumab as Initial Treatment of Advanced or Metastatic RET Fusion-Positive Non-Small Cell Lung Cancer	NCT04194944	Participants with RET fusion-positive non-squamous NSCLC that has spread to other parts of the body	Selpercatinib; carboplatin; cisplatin; pemetrexed; pembrolizumab	Pemetrexed with or without pembrolizumab	The primary endpoint was PFS. The secondary endpoints were ORR, DCR, PFS, OS, DOR and Time to Deterioration of Pulmonary Symptoms
A Randomized, Double-Blind, Phase III Study of Carboplatin-Paclitaxel/Nab-Paclitaxel Chemotherapy With or Without Pembrolizumab (MK-3475) in First Line Metastatic Squamous Non-small Cell Lung Cancer Subjects (KEYNOTE-407)	NCT03875092	Chinese adults with first line metastatic squamous NSCLC	Pembrolizumab; paclitaxel; nab-paclitaxel; carboplatin	Chemotherapy	The primary endpoints were PFS and OS. The secondary endpoints were ORR, DOR and AE
A Phase 3 Study of Pembrolizumab in Combination With Carboplatin/Taxane (Paclitaxel or Nab-paclitaxel) Followed by Pembrolizumab With or Without Maintenance Olaparib in the First-line Treatment of Metastatic Squamous Non-small Cell Lung Cancer (NSCLC)	NCT03976362	Have not received prior systemic treatment for their advanced/metastatic NSCLC	Pembrolizumab; carboplatin; paclitaxel; nab-paclitaxel; olaparib	Pembrolizumab + carboplatin + taxane + olaparib placebo	The primary endpoints were PFS and OS. The secondary endpoints were AE, EORTC QLQ-C30/LC13 score and TTD in EORTC QLQ-C30/LC13
PD-1 Inhibitors and Chemotherapy With Concurrent Irradiation at Varied Tumour Sites in Advanced Non-small Cell Lung Cancer	NCT03774732	Patients with advanced (stage IIIB/IV) NSCLC	Radiotherapy; pembrolizumab; chemotherapy	Pembrolizumab + chemotherapy	The primary endpoint was OS. The secondary endpoints were Tumour response, PFS, Acute/Late toxicities and Quality of life of the patients using EORTC-QLQ-C30
A Phase 3 Study of Pembrolizumab (MK-3475) in Combination With Concurrent Chemoradiation Therapy Followed by Pembrolizumab With or Without Olaparib vs. Concurrent Chemoradiation Therapy Followed by Durvalumab in Participants With Unresectable, Locally Advanced, Stage III Non-Small Cell Lung Cancer (NSCLC)	NCT04380636	Participants with unresectable, locally advanced NSCLC	Pembrolizumab; olaparib; etoposide; carboplatin; cisplatin; paclitaxel; pemetrexed; thoracic radiotherapy; durvalumab	Chemoradiation→durvalumab	The primary endpoints were PFS and OS. The secondary endpoints were AE, DOR, ORR and EORTC-QLQ-C30 score
A Phase 3 Study of Pembrolizumab in Combination With Pemetrexed/Platinum (Carboplatin or Cisplatin) Followed by Pembrolizumab and Maintenance Olaparib vs. Maintenance Pemetrexed in the First-Line Treatment of Participants With Metastatic Nonsquamous Non-Small-Cell Lung Cancer	NCT03976323	Advanced NSCLC with no EGFR mutation, ALK translocation or ROS1-directed therapy	Pembrolizumab; pemetrexed; carboplatin; cisplatin; olaparib	Pembrolizumab + pemetrexed + platinum therapy + pemetrexed	The primary endpoints were PFS and OS. The secondary endpoints were AE, EORTC-QLQ-C30 score and TTD in EORTC QLQ-LC13 Cough (Item 31) Scale Score
A Randomized Trial of Consolidative Immunotherapy With vs. Without Thoracic Radiotherapy and / or Stereotactic Body Radiation Therapy (SBRT) After First-Line Systemic Therapy for Metastatic NSCLC	NCT03867175	Stage IV NSCLC with no EGFR mutation, ROS1 or ALK gene rearrangements	Stereotactic body radiation therapy; pembrolizumab	Body radiation therapy	The primary endpoint was PFS. The secondary endpoints were Time of Progression, AE, Rate of Failure and Number of Participants with New Sites of Disease
A Randomized, Phase 3, Open-Label Study of Combinations of REGN2810 (Anti-PD-1 Antibody), Platinum-based Doublet Chemotherapy, and Ipilimumab (Anti-CTLA-4 Antibody) Versus Pembrolizumab Monotherapy in First-Line Treatment of Patients With Advanced or Metastatic Non-Small Cell Lung Cancer With Tumors Expressing PD-L1 $\geq$ 50%	NCT03515629	Patients with advanced squamous or non-squamous NSCLC whose tumors express PD-L1 in $\geq$ 50% of tumor cells	REGN2810/ipi; REGN2810/chemo/ ipi; Pembrolizumab	Pembrolizumab	The primary endpoint was PFS. The secondary endpoints were OS, ORR, TEAEs, SAEs, DLTs and Quality of life
QUILT 2.023: A Phase 3, Open-Label, 3-Cohort Randomized Study of N-803, in Combination With Current Standard of Care vs. Standard of Care as First-Line Treatment for Patients With Advanced or Metastatic NSCLC.	NCT03520686	Stage III/IV advanced or metastatic NSCLC	N-803; pembrolizumab; carboplatin; nab-paclitaxel	Control A: pembrolizumab; Control B: carboplatin + nab-paclitaxel or paclitaxel + pembrolizumab; control c: cisplatin or carboplatin + pembrolizumab + pemetrexed	The primary endpoint was PFS. The secondary endpoints were OS, ORR, PFS, DCR and Quality of life
A Randomized, Open-Label, Phase 3 Study of Pralsetinib Versus Standard of Care for First Line Treatment of RET Fusion-positive, Metastatic Non-Small Cell Lung Cancer	NCT04222972	Patients with RET fusion-positive metastatic NSCLC who have not previously received systemic anticancer therapy for metastatic disease	Pralsetinib; carboplatin; cisplatin; pemetrexed; pembrolizumab; gemcitabine	Platinum doublet with or without pembrolizumab	The primary endpoint was PFS. The secondary endpoints were OS, ORR, AE, DCR and CBR
A Phase 3, Randomized, Double-Blind, Placebo-Controlled, Multicenter Study Comparing Niraparib Plus Pembrolizumab Versus Placebo Plus Pembrolizumab as Maintenance Therapy in Participants Whose Disease Has Remained Stable or Responded to First-Line Platinum Based Chemotherapy With Pembrolizumab for Stage IIIB or IV Non-Small Cell Lung Cancer	NCT04475939	Participants with Stage IIIB or IV NSCLC who have achieved Stable disease (SD), Partial response (PR), or complete response (CR) following completion of platinum-based first-line induction chemotherapy with pembrolizumab	Niraparib; pembrolizumab	Placebo + pembrolizumab	The primary endpoints were PFS and OS. The secondary endpoints were AE, TTD in Lung Symptoms and TPP
A Randomized Non-inferiority Trial Evaluating the Length of Treatment With PD-1/PD-L1 Inhibitors in Patients With Advanced Solid Tumors	NCT04157985	Patients who have disease stability to stop treatment at 1 year or continue treatment until disease progression	Pembrolizumab; nivolumab; atezolizumab; ipilimumab	Continue Treatment with PD-1/PD-L1 inhibitor	The primary endpoint was Time to next treatment. The secondary endpoints were OS, irAEs and BOR
An Open-label, Randomized Phase III Study of Early Switch Maintenance vs. DElayed Second-line Nivolumab in Advanced Stage Squamous Non Small-cell Lung Cancer (NSCLC) Patients After Standard First-line Platinum-based Chemotherapy - EDEN Trial	NCT03542461	Patients with advanced stage squamous NSCLC without immunotherapy	Nivolumab	Best supportive care	The primary endpoint was OS. The secondary endpoints were PFS, TTF, OSind and PFSind
A Phase 3, Randomized, Open Label Study to Compare Nivolumab Plus Concurrent Chemoradiotherapy (CCRT) Followed by Nivolumab Plus Ipilimumab or Nivolumab Plus CCRT Followed by Nivolumab vs. CCRT Followed by Durvalumab in Previously Untreated, Locally Advanced Non-small Cell Lung Cancer (LA NSCLC)	NCT04026412	Participants with untreated locally advanced NSCLC	Nivolumab; ipilimumab; durvalumab	Nivolumab/durvalumab + CCRT	The primary endpoints were PFS and OS. The secondary endpoints were AE, ORR and TTR
A Phase IIIB/IV Safety Trial of Nivolumab (BMS-936558) in Subjects With Advanced or Metastatic Non-Small Cell Lung Cancer Who Have Progressed During or After Receiving At Least One Prior Systemic Regimen	NCT02066636	Advanced or metastatic NSCLC (subjects with non-squamous histology must be tested for EGFR mutations and ALK rearrangement)	Nivolumab		The primary endpoint was AEs. The secondary endpoint were AEs and Median time to onset and median time to resolution
An Open-Label, Randomized Phase 3 Trial of Nivolumab, or Nivolumab Plus Ipilimumab, or Nivolumab Plus Platinum Doublet Chemotherapy Versus Platinum Doublet Chemotherapy in Subjects With Chemotherapy-Naïve Stage IV or Recurrent Non-Small Cell Lung Cancer (NSCLC)	NCT02477826	Stage IV or recurrent NSCLC, with no prior systemic anticancer therapy	Nivolumab; ipilimumab; carboplatin; cisplatin; gemcitabine; pemetrexed; paclitaxel	Platinum doublet chemotherapy	The primary endpoints were PFS and OS. The secondary endpoints were ORR and LCSS
Open-Label, Randomized Trial of Nivolumab (BMS-936558) Plus Pemetrexed/Platinum or Nivolumab Plus Ipilimumab (BMS-734016) vs. Pemetrexed Plus Platinum in Stage IV or Recurrent Non-Small Cell Lung Cancer (NSCLC) Subjects With Epidermal Growth Factor Receptor (EGFR) Mutation Who Failed 1L or 2L EGFR Tyrosine Kinase Inhibitor Therapy	NCT02864251	Patients with EGFR mutation, NSCLC who failed first line (1L) or second-line (2L) EGFR TKI therapy	Nivolumab; ipilimumab; pemetrexed; cisplatin; carboplatin	Platinum doublet chemotherapy	The primary endpoint was PFS. The secondary endpoints were PFS, ORR, OS and DOR
Randomized, OpenLabel, Phase 3 Trial of Nivolumab Plus Ipilimumab or Nivolumab Plus Platinum Doublet Chemotherapy Versus Platinum Doublet Chemotherapy in Early Stage NSCLC	NCT02998528	Early stage IB-IIIa, operable NSCLC	Nivolumab; cisplatin; vinorelbine; gemcitabine; docetaxel; pemetrexed; carboplatin; paclitaxel; ipilimumab	Platinum doublet chemotherapy	The primary endpoints were EFS and pCR. The secondary endpoints were OS, MPR and TTDM
An Open-Label Randomized Phase III Trial of BMS-936558 (Nivolumab) Versus Docetaxel in Previously Treated Advanced or Metastatic Squamous Cell Non-small Cell Lung Cancer (NSCLC)	NCT01642004	Subjects with squamous cell NSCLC, after failure of prior platinum-based chemotherapy	Nivolumab; docetaxel	Docetaxel	The primary endpoint was OS. The secondary endpoints were PFS, ORR, OS and TTR
Randomized Phase III Study Testing Nivolumab and Ipilimumab Versus a Carboplatin Based Doublet in First Line Treatment of PS 2 or Elderly (More Than 70 Years Old) Patients With Advanced Non-small Cell Lung Cancer	NCT03351361	Patients already have metastatic disease and a systemic, palliative treatment is the primary therapeutic option	Nivolumab; ipilimumab; chemotherapy	Chemotherapy	The primary endpoint was OS. The secondary endpoints were PFS, ORR, Quality of life score and PD-L1
An Open Label, Safety Study of Participants With Non-Small Cell Lung Cancer Receiving Second-Line Nivolumab Monotherapy in Asia	NCT03195491	Patients in Asia with NSCLC who are treated with Nivolumab monotherapy as a second line or third line treatment	Nivolumab		The primary endpoint was AE. The secondary endpoints were AE and Laboratory test abnormalities
A Randomized Phase 3 Study of Sitravatinib in Combination With Nivolumab Versus Docetaxel in Patients With Advanced Non-Squamous Non-Small Cell Lung Cancer With Disease Progression On or After Platinum-Based Chemotherapy and Checkpoint Inhibitor Therapy SAPHIRE	NCT03906071	Patients with advanced non-squamous NSCLC who have previously experienced disease progression on or after platinum-based chemotherapy and checkpoint inhibitor therapy	Nivolumab; sitravatinib; docetaxel	Docetaxel	The primary endpoint was OS. The secondary endpoints were AEs, PFS and ORR
A Randomized Phase 3 Trial Comparing Continuation Nivolumab-Ipilimumab Doublet Immunotherapy Until Progression Versus Observation in Treatment-naïve Patients With PDL1-positive Stage IV Non-Small Cell Lung Cancer (NSCLC) After Nivolumab-Ipilimumab Induction Treatment	NCT03469960	Stage IV NSCLC, with no prior systemic anticancer therapy, PD-L1 tumor content $\geq$ 1% and <50%	Ipilimumab; nivolumab	Standard treatment: 6 months of treatment by nivolumab + ipilimumab then nivolumab + ipilimumab then in case of progression platinum-based doublet recommended	The primary endpoint was PFS. The secondary endpoints were OS, PFS and Quality of life
Adjuvant Nivolumab in Resected Lung Cancers (ANVIL) - A Randomized Phase III Study of Nivolumab After Surgical Resection and Adjuvant Chemotherapy in Non-Small Cell Lung Cancers	NCT02595944	Patients with stage IB-IIIa NSCLC	Nivolumab	No intervention	The primary endpoints were DFS and OS. The secondary endpoint was AE
Randomized Phase III Trial of Local Consolidation Therapy (LCT) After Nivolumab and Ipilimumab for Immunotherapy-Naïve Patients With Metastatic Non-Small Cell Lung Cancer (LONESTAR) -Strategic Alliance: BMS	NCT03391869	Patients with stage IV NSCLC	Radiation therapy; local consolidation therapy; nivolumab; ipilimumab	Ipilimumab + nivolumab	The primary endpoint was OS. The secondary endpoints were PFS, TANM and Quality of life
A Phase III Clinical Trial of Adjuvant Chemotherapy vs. Chemoimmunotherapy for Stage IB-IIIa Completely Resected Non-small Cell Lung Cancer (NSCLC) Patients.	NCT04564157	Stage IB-IIIa, completely resected, NSCLC patients	Carboplatin; paclitaxel; nivolumab	Carboplatin + paclitaxel	The primary endpoint was DFS. The secondary endpoints were OS and AE
A Phase 3, Randomized, Double-blind Study of Neoadjuvant Chemotherapy Plus Nivolumab Versus Neoadjuvant Chemotherapy Plus Placebo, Followed by Surgical Resection and Adjuvant Treatment With Nivolumab or Placebo for Participants With Resectable Stage II-IIIB Non-small Cell Lung Cancer	NCT04025879	Early-stage NSCLC with no prior systemic anti-cancer treatment	Nivolumab; carboplatin; cisplatin; paclitaxel; pemetrexed; docetaxel	Neoadj. Plac. + Pt-based doublet chemo followed by Adj.Plac.	The primary endpoint was EFS. The secondary endpoints were OS, pCR, MPR and AE
A Multicenter, Randomized, Double-Blind Trial in Subjects With Non-Squamous Non-Small Cell Lung Cancer (TASUKI-52)	NCT03117049	Stage IIIB/IV or recurrent non-squamous NSCLC unsuitable for radical radiation	ONO-4538; carboplatin; paclitaxel; bevacizumab	Placebo group	The primary endpoint was PFS. The secondary endpoints were OS, PFS, ORR and AE